
Code of Practice for Timber Production 2014 (as amended 2022)

Schedule 1: Management Standards and Procedures for timber harvesting operations in Victoria's State forests



Cover photo

Manna Gum (Eucalyptus viminalis) stand in wet eucalypt forest, Dandenong Ranges, Victoria

Acknowledgment

We acknowledge and respect Victorian Traditional Owners as the original custodians of Victoria's land and waters, their unique ability to care for Country and deep spiritual connection to it. We honour Elders past and present whose knowledge and wisdom has ensured the continuation of culture and traditional practices.

We are committed to genuinely partner, and meaningfully engage, with Victoria's Traditional Owners and Aboriginal communities to support the protection of Country, the maintenance of spiritual and cultural practices and their broader aspirations in the 21st century and beyond.



© The State of Victoria Department of Environment, Land, Water and Planning 2022

This work is licensed under a Creative Commons Attribution 4.0 International licence. You are free to re-use the work under that licence, on the condition that you credit the State of Victoria as author. The licence does not apply to any images, photographs or branding, including the Victorian Coat of Arms, the Victorian Government logo and the Department of Environment, Land, Water and Planning (DELWP) logo. To view a copy of this licence, visit creativecommons.org/licenses/by/4.0/

ISBN 978-1-76105-993-3 (pdf/online/MS word)

Disclaimer

This publication may be of assistance to you but the State of Victoria and its employees do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in this publication.

Accessibility

If you would like to receive this publication in an alternative format, please telephone the DELWP Customer Service Centre on 136 186, or email customer.service@delwp.vic.gov.au or via the National Relay Service on 133 677, www.relayservice.com.au. This document is also available on the internet at www.delwp.vic.gov.au

Contents

Co	Contents		
1.	Intr	oduction	6
	1.1	Scope	6
	1.2	Role	6
	1.3	Application	6
	1.4	Procedure for seeking exemptions or temporary variations	6
2.	Pla	nning and Record Keeping	7
	2.1	FMZ and Planning Information	7
	2.2	Communication and engagement	7
	2.3	Traditional owner rights and interests	8
	2.4	Forest Coupe Plans	8
	2.5	Coupe Size and configuration	10
3.	Wa	ter Quality, River Health and Soil Protection	11
	3.1	Waterway classification	11
	3.2	Soil Erosion and Water Quality Risk Assessment	11
	3.3	Waterway Protection	12
	3.4	Slope	13
	3.5	Water supply protection areas	13
4.	Bio	diversity	15
	4.1	Habitat Retention	15
	4.2	Fauna and Flora	18
	4.3	Vegetation communities	19
	4.4	Pests, weeds and diseases	23
5.	lmp	ortant values	24
	5.1	Heritage, historic places and Aboriginal cultural heritage	24
	5.2	Landscape Sensitivity Areas	24
	5.3	Apiary	27
	5.4	Recreation	27
	5.5	Research	28
	5.6	Fuel Hazard ratings in Fire Management Zones	28
6.	Roa	ading	32
	6.1	Road planning	32
	6.2	Road construction	33

	6.3	Road Maintenance	35
	6.4	Road Rehabilitation	35
7.	Timl	per Harvesting	36
	7.1	Coupe Management	36
	7.2	Coupe Infrastructure	37
	7.3	Operational restrictions	39
8.	Fire	Salvage Harvesting	40
9.	Fore	est Regeneration and Management	43
	9.1	Regeneration	43
	9.2	Thinning	45
Арр	endix	1: Tables and Figures	46

Glossary

All **bold** terms included in Schedule 1 are defined within the **Code** Glossary.

Acronyms

ANU The Australian National University

ASL above sea level

B **buffer**

CAR comprehensive, adequate and representative reserve system

cm centimetres

DBHOB **diameter** at breast height (1.3m) over bark

DC designated catchment identified in a **forest** management plan

EVC ecological vegetation class

F filter or filter strips

FMA forest management area

FMZ forest management zone

GMZ general management zone

GRU geographic representation unit

ha hectares

HEMS high elevation mixed species forest

km kilometres

LFP Long Footed Potoroo

m metres mm millimetres

RFA regional forest agreement

SAP special area plan

SMZ special management zone
SPZ special protection zone

SWSCA special water supply catchment area

1. Introduction

1.1 Scope

1.1.1.1 As part of the **Code**, The **Management Standards and Procedures** apply to all commercial **timber harvesting operations** conducted in Victoria's **State forests**.

1.2 Role

- 1.2.1.1 This Schedule provides standards and procedures to instruct managing authorities, harvesting entities and **operators** in interpreting the requirements set out in the main body of the **Code**.
- 1.2.1.2 These **Management Standards and Procedures** are in addition to the mandatory actions set out in the main body of the **Code**.

1.3 Application

1.3.1.1 Requests for exemptions or temporary variations to these **Management Standards and Procedures** will demonstrate to the satisfaction of the **Minister** or **delegate** that they are consistent with the Operational Goals and Mandatory Actions of the **Code**.

1.4 Procedure for seeking exemptions or temporary variations

- 1.4.1.1 If a provision in these **Management Standards and Procedures** provides for an exemption or temporary variation, the **managing authority** may submit an application form and any other required or supporting information to the **Minister**.
- 1.4.1.2 The **Minister** or **delegate** will consider the request, seeking additional information where required, and respond by providing a signed approval to the submitter.
- 1.4.1.3 Where the request is not **approved**, the response will include a justification of the decision.
- 1.4.1.4 The **Minister** or **delegate** will maintain a register of the outcomes for all exemption or temporary variation requests.

2. Planning and Record Keeping

2.1 FMZ and Planning Information

- 2.1.1.1 When planning **timber harvesting operations**, for information regarding the **FMZS** and locations of **Special Management Zones** (**SMZ**) or **Special Protection Zones** (**SPZ**), the **managing authority** will refer to the **FMZ**100 geospatial layer published on the Victorian Spatial Data Library.
- 2.1.1.2 Where an **SPZ**, a protection area or an **exclusion** area located within an **SMZ** is based on a modelled value or incorrectly mapped geographic feature and is determined not to exist in the field or is incorrectly located, the **managing authority** may notify the **Secretary** (or **delegate**) prior to commencement of the **timber harvesting operation** requesting an amendment to the boundary of the **SPZ**, protection area or **exclusion** area.

2.2 Communication and engagement

- 2.2.1.1 The **managing authority** must make maps and schedules of **coupes** that have been selected for **timber harvesting operations** and associated access roading available for public scrutiny prior to commencement of the operations. This information must provide:
 - a) clear maps showing the location of **coupes** and major access roading (including extensions or upgrading of the permanent road network); and
 - b) approximate timing and duration of timber harvesting operation schedules.
- 2.2.1.2 Consult the following parties during the establishment or amendment of plans, allowing adequate time to ensure interests are considered and operational arrangements are resolved prior to commencement of operations:
 - a) relevant bee keepers or bee keeping associations;
 - b) relevant **Department** regions regarding roading, burning and other relevant land management activities;
 - c) relevant water authorities when applying chemicals in a **special water supply catchment area**: and
 - d) relevant water supply authorities when **coupes** are proposed to occur within **special water supply catchment areas** in the Benalla-Mansfield **FMA** and North East **FMAs**; and
 - e) Melbourne Water regarding any **timber harvesting operations** which have potential to impact on the Water Resource Values within the Bunyip River, Tarago River, Thomson River and Yarra Tributaries **State forests**.

2.3 Traditional owner rights and interests

- 2.3.1.1 During planning for **timber harvesting operations**, have regard to obligations under the following:
 - i. the *Native Title Act 1993*, including any **Indigenous Land Use Agreement** made under that Act;
 - ii. the *Traditional Owner Settlement Act 2010*, including any **Recognition and Settlement Agreements**, **Land Use Activity Agreements** and **Natural Resource Agreements**;
 - iii. **Aboriginal cultural rights** under the *Charter of Human Rights and Responsibilities Act 2006.*

2.4 Forest Coupe Plans

- 2.4.1.1 Forest Coupe Plans prepared for timber harvesting operations must:
 - a) state the **net coupe area** that is planned to be harvested or tended;
 - b) state the start date and season during which operations are scheduled to occur;
 - c) identify the silvicultural systems or **tending** methods to be employed;
 - d) identify requirements for the location*, design, construction, maintenance and use of **roads**, and closure of **temporary roads**;
 - e) identify requirements for the design, siting*, construction, use, and rehabilitation of log landings and dumps and, where necessary, siting and rehabilitation measures for major snig tracks;
 - f) describe **regeneration** procedures to be applied;
 - g) identify and describe management actions for all relevant coupe values specified within the Code and the FMZS;
 - h) describe measures employed to protect **biodiversity** such as:
 - iv. habitat tree retention;
 - v. provision for the continuity and replacement of old **hollow bearing trees** within the harvestable area; and
 - vi. retention of long-lived **understorey** species in appropriate numbers and configurations to allow for the **regeneration** of the vegetation types that existed within the **coupe** before the **timber harvesting operation** commenced.
 - i) describe soil erosion hazard classes present within the coupe and measures employed to protect and rehabilitate soils and to maintain water quality and river health.

^{*} The location of **temporary roads** and **coupe infrastructure** may be specified in the field and added subsequently to the commencement of operations.

- 2.4.1.2 Forest Coupe Plans must include a map which clearly and accurately identifies:
 - a) the **net coupe area** that is planned to be harvested;
 - areas within and adjacent to a coupe that are to be excluded from timber harvesting
 operations and associated activities, or to which special prescriptions apply (including
 biodiversity protection or habitat enhancement, water quality and aquatic habitat protection,
 landscape protection, historic places and Aboriginal cultural heritage); and
 - c) waterway, historic tramway and water race crossing locations.
- 2.4.1.3 Forest Coupe Plans prepared for road construction or significant road improvement operations also identify:
 - a) the class of road to be constructed / improved;
 - b) maximum clearing width;
 - c) sections of the **road** subject to narrowed or increased easements;
 - d) methods and location of proposed crossings and drainage structures; and
 - e) the **FMZs** impacted upon and any requirements relevant to approvals associated with roading operations within **SMZ** or **management areas** as well as **SPZ** or **protection areas**.

2.5.1 Clearfall, seed tree or shelterwood 1 coupes 2.5.1.1 The maximum size of clearfall, seed tree or shelterwood 1 coupes is 40 ha net harvested area. 2.5.1.2 The maximum aggregated size of coupes that contain clearfall, seed tree or shelterwood 1 is 120 ha **net harvested area** within a 5 year period. 2.5.2 **Shelterwood 2 coupes** 2.5.2.1 Shelterwood 2 coupes may exceed 40 ha net harvested area if adequately stocked with eucalypts that are 5 or more years old. 2.5.3 Thinning coupes 2.5.3.1 The maximum size of a **thinning coupe** is 120 ha **net harvested area**. 2.5.4 Single tree selection coupes 2.5.4.1 Single tree selection coupes may be of any size. 2.5.5 **Uneven-aged coupes** 2.5.5.1 The maximum size of an uneven-aged coupe is 120 ha net harvested area. 2.5.5.2 No gaps or patches created by seed tree harvesting operations are to be more than 20 ha in area. 2.5.5.3 The cumulative area of gaps between 1 and 20 ha must comprise no more than 33 % of the net harvested area. 2.5.5.4 More than 66 % of the net harvested area must comprise areas stocked by basal area retention with the balance stocked with regeneration. 2.5.6 Salvage coupes 2.5.6.1 Salvage coupes may exceed standard area limits in accordance with the fire salvage harvesting area limits of Clause 8.1.3.1. 2.5.7 All coupe types 2.5.7.1 20 m is the minimum width of vegetation to be retained between **coupe** aggregates that have both been harvested within the last 5 years. 2.5.7.2 Adjacent partially stocked or unstocked areas may be incorporated into the coupe for regeneration provided that the coupe is within the coupe size limit.

2.5

Coupe Size and configuration

3. Water Quality, River Health and Soil Protection

3.1 Waterway classification

- 3.1.1.1 Use the following categories when determining **buffer** (B) and **filter** (F) widths for **waterways** within and immediately adjacent to each **coupe**. Aids to the identification of each class of **waterway** are provided in the **Code** Glossary.
 - a) Permanent streams, pools and wetlands.
 - b) Temporary streams.
 - c) Drainage lines.

3.2 Soil Erosion and Water Quality Risk Assessment

3.2.1 Statewide

- 3.2.1.1 Conduct field assessments to determine the soil erosion hazard and soil **permeability** classifications for an area proposed for any soil disturbing **timber harvesting operations** as follows.
- 3.2.1.2 Collect soil profile samples that reflect the variety of soils represented within the coupe.
 Note: Significant changes in forest structure and composition often indicate a variation in soil type.
- 3.2.1.3 Score each soil factor using **Table 1 Soil susceptibility to breakdown** and **Table 2 Soil permeability** for each horizon within the soil profile to the anticipated depth of disturbance.
- 3.2.1.4 Calculate the soil susceptibility to breakdown rating for horizons A and B (and C if likely to be disturbed in operations) using **Table 1 Soil susceptibility to breakdown**.
- 3.2.1.5 Calculate the soil permeability rating for horizons A and B (and C if likely to be disturbed in operations) using **Table 2 Soil permeability**.
- 3.2.1.6 Calculate the soil erosion site factor rating for the site using **Table 3 Soil erosion site factor**. Use **Figure 1 Rainfall erosivity** contours in non-SI units for Victoria to determine the erosivity index.
- 3.2.1.7 Calculate the **soil erodibility** classification for horizons A and B (and C if likely to be disturbed in operations) using **Table 4 Soil erodibility classification**.
- 3.2.1.8 Calculate the soil erosion hazard classification for horizons A and B (and C if likely to be disturbed in operations) using **Table 5 Soil erosion hazard classification**.
- 3.2.1.9 Calculate the soil **permeability** classification for horizons A and B (and C if likely to be disturbed in operations) using

3.2.1.10

3.2.1.11 Table 6 Soil permeability classification.

- 3.2.1.12 Calculate the overall soil category for horizons A and B (and C if likely to be disturbed in operations) using
- 3.2.1.13
- 3.2.1.14 **Table 7 Overall soil category**.
- 3.2.1.15 Use the overall soil category for the horizon with the highest category to determine water quality risk as shown in **Table 8 Water quality risk**.

3.3 Waterway Protection

3.3.1 Statewide

3.3.1.1 Apply the appropriate protection to class of waterway as outlined in Table 9 Minimum widths in metres for buffers (B) and filter (F) strips applicable to various waterway categories, in relation to water quality risk and slope.

Notes:

- Slope is the average slope of the **coupe** area in the vicinity of the water body.
- **Buffers** and **filter strips** must be applied to each **waterway** class regardless of the origins of the channelling.
- The width of **buffer** areas and **filter strips** must be measured in the horizontal plane, from the edge of the saturated zone (at time of harvesting) or channel (whichever is greater), on each side of the **waterway**.

3.3.2 Bendigo FMA

- 3.3.2.1 **Drainage lines** are considered to be present if banks and/or eroded sections greater than 30 cm deep and greater than 10 m long exist. Distinctive **riparian vegetation** such as River Red Gum (*Eucalyptus camaldulensis*) will generally be absent. Where sections meeting these criteria occur within 50 m, the interconnecting depression is treated as part of the **drainage line**.
- 3.3.2.2 Machinery movement within 10 m of depressions occurring above **drainage lines** will be directed along the contour.

3.3.3 Mid Murray FMA

- 3.3.3.1 Exclude **timber harvesting operations** within 60 m from the high bank of the Murray River.
- 3.3.3.2 Exclude **timber harvesting operations** within 40 m from the high bank of the Gunbower Creek and the Ovens River.
- 3.3.3.3 Apply a 20 m **buffer** to main rivers, anabranches, lakes, billabongs and lagoons that maintain permanent open water at minimum river flows.
- 3.3.3.4 When water is present, a 10 m **buffer** applies to **temporary effluent** and/or **confluent streams**.
- 3.3.3.5 When water is absent, a 10 m filter applies to temporary effluent and/or confluent streams.
- 3.3.3.6 **Temporary effluent** and/or **confluent streams** have sharply defined channels, relatively deep flowing water and overland flow in the immediate vicinity during flood.
- 3.3.3.7 If distinctive **riparian vegetation** is present and wider than 3.3.3.1, **buffers** and **filters** are measured from the edge of the distinctive **riparian vegetation**.

3.4 Slope

- 3.4.1 Slope restrictions
- 3.4.1.1 Exclude **timber harvesting operations** from slopes over 30 degrees.
- 3.4.1.2 Exclude **timber harvesting operations** from slopes over 25 degrees in the East Gippsland **FMA** in areas with granite-based soils.
- 3.4.1.3 Up to 10% of the planned **net coupe area** can contain areas greater than the slope limits referred to in 3.4.1.1, 3.4.1.2, 3.5.1.1 and **Table 11 Water supply protection areas**, where the risk of mass soil movement has been managed accordingly.

3.5 Water supply protection areas

- 3.5.1.1 Apply the slope limits, seasonal closures, **buffer** and **filter strip** widths and other relevant management actions specified in **Table 11 Water supply protection areas** for **timber harvesting operations** and associated roading and **regeneration** in **water supply protection areas**.
- Refer to **Table 11 Water supply protection areas** for actions that apply to **water supply protection area SMZ**s in the Benalla-Mansfield **FMA**, the East Gippsland **FMA** and Midlands **FMA**.
- 3.5.1.3 Where any proposed **timber harvesting operations** are not in accordance with clauses 3.5.1.1 or 3.5.1.2, obtain exemption approval from the **Minister** or **delegate** in accordance with Section 1.4.
- 3.5.1.4 Special water supply catchment areas and water supply protection areas not listed in Table 11 Water supply protection areas do not require protection in addition to existing Code requirements.
- 3.5.1.5 In addition to **Table 11 Water supply protection areas**, in the Bunyip, Thomson and Tarago **special water supply catchment areas** and the Yarra Tributaries **State forests** the area harvested must not exceed the following limits measured as a 10-year rolling average:
 - a) Thomson Ash forests 150 ha/year, mixed species forests 15ha/year;
 - b) Tarago Ash forests 55 ha/year, mixed species forests 23ha/year;
 - c) Yarra Tributaries Ash forests 52 ha/year, mixed species forests 15 ha/year; and
 - d) Bunyip Ash forests 15 ha/year, mixed species forests 15 ha/year.
- 3.5.1.6 In addition to **Table 11 Water supply protection areas**, harvesting in the Learmonths Creek special water supply catchment must not be undertaken if it would cause:
 - a) harvesting of Ash forest in the catchment to exceed an average harvest of 7 ha per annum, which average is to be calculated progressively in each financial year (including as at 30 June in each financial year) using the area of Ash forest harvested in that financial year and the areas harvested in each of the preceding nine financial years; or
 - b) harvesting of **mixed species forest** in the catchment to exceed an average of 3 ha per annum, which average is to be calculated progressively in each financial year (including as at 30 June in each financial year) using the area of **mixed species forest** harvested in that financial year and the areas harvested in each of the preceding nine financial years.
- 3.5.1.7 In addition to **Table 11 Water supply protection areas**, in the Benalla-Mansfield **FMA**, the total area of **coupes** harvested using **even-aged** harvesting and **regeneration** systems, such

as **clearfall** or **seed tree harvesting** operations, must not exceed 5% of the area of public land in designated catchments in any 3-year period.

3.5.1.8 In addition to **Table 11 Water supply protection areas**, in the Rocky and Betka River catchments in the East Gippsland **FMA**, no new **road** crossings may be built on major streams.

4. Biodiversity

c) where they exist:

4.1	Habitat Retention
4.1.1	Statewide
4.1.1.1	Retain habitat trees in accordance with the relevant FMA summary provided in Table 12 Habitat tree prescriptions.
4.1.1.2	Trees retained in buffers or exclusion areas that have been extended beyond minimum required widths or radius can contribute to habitat tree retention requirements.
4.1.1.3	Retain all living large trees and protect them from the direct effects of timber harvesting operations and regeneration burning. This includes not deliberately felling, falling into or damaging large trees during harvesting and extraction, and ensuring slash is not accumulated within 3m of the base of large trees immediately prior to regeneration burns.
4.1.1.4	Where possible, incorporate large trees in retained patches of habitat trees or expanded buffers or exclusion areas .
4.1.1.5	Include all large trees in a Large Tree Register.
4.1.2	Benalla/Mansfield FMA and North East FMA
4.1.2.1	When selecting habitat trees , prioritise live hollow bearing trees where they are present and trees of younger age classes that are likely to develop hollows in the longer term.
4.1.2.2	Where possible retain dead trees for habitat trees .
4.1.2.3	Retain habitat trees adjacent to areas of high value habitat and areas most easily protected from damage during harvesting and regeneration operations.
4.1.2.4	Habitat trees may be retained in groups/patches dispersed across the coupe.
4.1.2.5	Seed trees may be counted as habitat trees.
4.1.3	Bendigo FMA and Box-Ironbark forests in the Midlands FMA.
4.1.3.1	Permanently retain:
	 a) all trees (both standing dead or living) greater than 20 cm DBHOB with visible hollows where safe and practicable to do so. These trees may count towards retention requirements in clause (b) and (c);
	b) at least 10 living trees per hectare between 30 cm and 39.9 cm DBHOB;

unmerchantable trees with healthy crowns should be preferentially retained;

ii). 2 living trees per hectare between 50 and 59.9 cm DBHOB inclusive.

i). 2 living trees per hectare between 40 cm and 49.9 cm DBHOB inclusive; and

d) all trees greater than 60 cm DBHOB in high quality sawlog harvesting operations;

- e) all trees greater than 40 cm DBHOB in low quality sawlog and firewood harvesting operations;
- f) within **management areas** or **SMZ**, all trees greater than 40 cm DBHOB in high and low quality sawlog and firewood harvesting operations;
- g) all standing dead trees greater than 40 cm DBHOB;
- h) all living Yellow Box, Fryers Range Scentbark and Bealiba Ironbark trees. These trees may count towards retention requirements.
- 4.1.3.2 A whole of **coupe** approach should be adopted (numbers averaged over **coupe** area) to ensure that the best **habitat trees** are retained, including where these occur in groups.
- 4.1.3.3 Firewood harvesting within a **management area** or **SMZ** must be managed to ensure coarse woody debris levels are not reduced below pre-harvest levels.
- 4.1.3.4 All log sections larger than 40cm **diameter** with **hollows** larger than 10cm **diameter** must be retained.
- 4.1.3.5 The period between the next and subsequent sawlog harvesting operations will be at least 25 years to allow recruitment across all habitat classes. **Habitat trees** intended for permanent retention will not be permanently tagged or labelled. It is expected that these trees will be evident from their form.

4.1.4 Central Highlands FMAs

- 4.1.4.1 When selecting **habitat trees**, prioritise **hollow bearing trees** where they are present and trees most likely to develop **hollow**s in the short term.
- 4.1.4.2 Scatter habitat trees across the timber harvesting coupe in mixed species forest.
- 4.1.4.3 Where possible, retain potential **hollow** bearing **ash** eucalypts in clumps to increase their protection from exposure, windthrow and fire.
- 4.1.4.4 No gap between retained vegetation is to be greater than 150m.
- 4.1.4.5 Retain **habitat trees** where they can be most easily protected from damage during **timber harvesting operations** and **site preparation** treatment.

4.1.5 East Gippsland FMA and Gippsland FMAs

- 4.1.5.1 When selecting **habitat trees**, prioritise old living trees with a range of **hollow** sizes. Where these are absent or not present in sufficient numbers, prioritise trees that are old enough to develop **hollows** during the next 50 years.
- 4.1.5.2 Stags and younger, smaller trees may be counted as **habitat trees** if trees of the type described in 4.1.5.1 are absent or not present in sufficient numbers.
- 4.1.5.3 Where possible, retain **habitat trees** in small clusters which include younger **regrowth** and **understorey**.
- 4.1.5.4 Distribute **habitat tree** clusters across the **coupe** with consideration of the proximity of other retained vegetation.

4.1.6 Horsham FMA

4.1.6.1 Retain trees that have signs that they are being actively utilised or occupied by **wildlife** as a priority over potential **habitat trees** and **dead** trees.

4.1.6.2 **Habitat trees**:

- a) are large actively growing trees with a spreading form; and/or
- b) have hollows present and forming.
- 4.1.6.3 Trees growing in ephemeral **wetlands** and within 20 m of the edge of the **wetland** may only be harvested using single tree selection. A 20 m **buffer** from the water line or saturated zone, wherever it occurs at the time of harvesting, also applies.
- 4.1.6.4 In areas of Red-tailed Black-Cockatoo potential habitat that is not protected by a **protection** area, SPZ, management area or SMZ, do not remove any hollow bearing trees, tees of the species Brown Stringybark or trees of the species Desert Stringybark with DBHOB > 45cm.
- 4.1.7 Midlands FMA, except the Box Ironbark forests
- 4.1.7.1 In the non **Box Ironbark forests**, the **habitat tree** numbers in **Table 12 Habitat tree prescriptions** are the maximum required.
- 4.1.7.2 Prioritise retention of **hollow bearing trees** or gum species as **habitat trees**.

4.1.8 Mid Murray FMA

- 4.1.8.1 **Habitat tree** requirements in **Table 12 Habitat tree prescriptions** are the minimum requirement.
- 4.1.8.2 When selecting **habitat trees**, prioritise **hollow bearing trees** where they are present and trees most likely to develop **hollows** in the short term.
- 4.1.8.3 In addition to **habitat tree** requirements in **Table 12 Habitat tree prescriptions**, retain all trees known to be used for nesting by significant **fauna** (e.g. Superb Parrot and Regent Parrot).
- 4.1.8.4 Retain all naturally fallen wood and any **dead** standing trees with **hollows**.

4.1.9 Otway FMA

- 4.1.9.1 Habitat tree numbers in Table 12 Habitat tree prescriptions are a minimum requirement.
- 4.1.9.2 Retain existing and potential **habitat trees** in regularly configured clusters or on the edges of **coupes**.
- 4.1.9.3 Design **coupe** boundaries and **habitat tree** clusters so there is no more than 200m between areas of retained habitat.
- 4.1.9.4 Where located on or adjacent to a **coupe** boundary, habitat patches are not included in the net harvest area of adjoining **coupes**.

4.1.10 Portland FMA

4.1.10.1 Habitat trees:

- a) are living;
- b) have a DBHOB greater than 30cm;
- c) are fully mature;
- d) have existing hollows (in a minimum of 50% of trees retained for habitat); and

- e) cater for identified key species needs.
- 4.1.10.2 Prioritise retention of **habitat trees**:
 - a) greater than 80cm diameter;
 - b) in clusters rather than scattered throughout the **coupe**;
 - c) located more than 20m of any road, track or coupe boundary likely to be used as a fire break; and
 - d) located in places that extend or link **filter strips**, streams or other retained areas.
- 4.1.10.3 Retain all **dead** trees if they contain **hollows** and are greater than 50cm DBHOB. **Dead** trees are not counted in the minimum **habitat tree** requirements.
- 4.1.10.4 In areas of Red-tailed Black-Cockatoo potential habitat that is not protected by **SPZ** or **SMZ**, do not remove any **hollow bearing trees**, tees of the species Brown Stringybark or trees of the species Desert Stringybark with DBHOB > 45cm.

4.2 Fauna and Flora

- 4.2.1 Detection-based management
- 4.2.1.1 Detection based management obligations apply in any area that may be affected by current or planned **timber harvesting operations**, and in any area in which an obligation may affect the conduct of such **timber harvesting operations** (for example, if a **protection area** would include an area within which **timber harvesting operations** are proposed).
- 4.2.1.2 If evidence of the presence of a value listed in Table 13 Rare or threatened fauna and invertebrate prescriptions or Table 14 Rare or threatened flora prescriptions is identified, the managing authority must:
 - a) notify the **Secretary**, providing details (including spatial information) of evidence and the value location; and
 - b) unless the **Secretary** otherwise approves, take appropriate steps to **verify** evidence of the presence of the value.

Note: The **Secretary** may otherwise approve if the **Secretary** intends to take steps to **verify** the existence of the value.

- 4.2.1.3 If evidence of the presence of a value listed in Table 13 Rare or threatened fauna and invertebrate prescriptions or Table 14 Rare or threatened flora prescriptions is verified, apply and undertake any associated management action specified in the Table.
- 4.2.1.4 If a management action or other mandatory action includes a requirement to apply a **protection** area or a **management area**, the **managing authority** must (in addition to any other action):
 - a) Provide information to the **Secretary** about the extent and boundary of the area proposed to be delineated as a **protection area** or **management area**;
 - b) Follow any input or direction of the **Secretary** about the extent and boundary of the area (which must not be inconsistent with the **Code**):
 - c) Delineate the boundary of the **protection area** or the **management area** in any applicable **Forest Coupe Plan** (consistently with the applicable requirement and any

- input or direction from the **Secretary**), including by updating the **Forest Coupe Plan** as the case requires; and
- d) Keep sufficient records of the boundaries of any part of an area for which there is not an applicable **Forest Coupe Plan** to ensure that the area is identifiable for the purpose of compliance with the **Code** and to enable that area to be delineated on any **Forest Coupe Plan** that subsequently includes that area.
- 4.2.1.5 In any **management area** that is required to be applied by these **Management Standards and Procedures**, the **managing authority** must (in addition to any other action):
 - a) consult with the **Secretary** about proposed measures to ensure the persistence of a value across the **management area**; and
 - b) ensure sufficient practical measures are undertaken to ensure the persistence of a value across the **management area**.
- 4.2.1.6 The **managing authority** must comply with the requirements of clause 4.2.1.2, 4.2.1.3, 4.2.1.4 and 4.2.1.5, either:
 - a) prior to the commencement of timber harvesting operations; or
 - b) if a requirement only arises after **timber harvesting operations** have commenced, as soon as possible after the requirement arises.

4.3 Vegetation communities

4.3.1 Box Ironbark

- 4.3.1.1 In the **Gippsland FMAs** exclude **selective harvesting** from **Box Ironbark forests** typically containing **Forest** Red Gum (*Eucalyptus tereticornis*), Yellow Box (*Eucalyptus melliodora*), Coast Grey Box (*Eucalyptus bosistoana*) and Red Ironbark (*Eucalyptus tricarpa*). Silvicultural practices that promote **regeneration** of these species is permitted.
- 4.3.1.2 In the East Gippsland **FMA** exclude **selective harvesting** from **Box Ironbark forests** typically containing Red Ironbark (*Eucalyptus tricarpa*), Gippsland Grey Box (*Eucalyptus bosistoana*), Red Box (*Eucalyptus polyanthemos*), Blue Box (*Eucalyptus baueriana*) and Yellow Stringybark (*Eucalyptus muelleriana*). The use of **seed tree harvesting** systems is permitted to restore the original species mix when combined with:
 - cutting stumps of desired species to a maximum height of 30 cm, to encourage coppice growth;
 - b) supplementary planting and sowing where necessary;
 - c) removing unproductive trees of the less-preferred species to remove **overwood** competition; and
 - d) thinning of advanced regrowth.

4.3.2 Heathland

- 4.3.2.1 Avoid **road construction** across areas of **heathland** or within 40m of **heathlands** unless no reasonable alternative exists.
- 4.3.2.2 In the **Gippsland FMAs**, exclude Wet **Heathland**, Clay **Heathland** and Riparian Scrub Mosaic **EVC**s from harvesting. Protect these **heathland EVC**s with a 40m **buffer**.
- 4.3.2.3 In the East Gippsland FMA and Otway FMA where evidence of heathland is found in the field and it isn't already classified as SPZ, apply a protection area prior to commencement of the timber harvesting operation consistent with management actions listed in Table 15 Detection based rules for Heathland, Snow Gum, Swamp Gum and Heath vegetation communities.
- 4.3.2.4 In the **Central Highlands** and **North East FMAs**, where evidence of **heathland** is found in the field and it isn't already classified as **SPZ**, apply a **protection area** over the identified **heathlands**, however there is no **buffer** required.

4.3.3 Montane Riparian Thicket (MRT)

- 4.3.3.1 In the Tambo **FMA** protect small **stands** of **MRT** between 0.01ha and 0.5ha and less than 10m wide with a 10m **filter strip** and **stands** of **MRT** wider than 10m with a 20m wide **filter strip**. Protect **stands** of **MRT** greater than 0.5ha with a 20m **buffer** from the edge of the Mountain Tea-tree **canopy**.
- 4.3.3.2 In all other **FMAs** apply the **heathland** prescriptions listed above in 4.3.2.
- 4.3.3.3 **MRT stands** must contain at least 40 % **canopy** cover of Mountain Tea-tree (*Leptospermum grandifolium*) and at least two key **understorey** species from the **MRT** definition in the **Glossary**. Small gaps in the Mountain Tea-tree **canopy** may occur at intervals up to 10m in length. Where **MRT stands** are separated by **canopy** gaps greater than 10m, they are to be treated as individual **stands**.

4.3.4 Old growth forest

- 4.3.4.1 Within the **Central Highland's FMAs**, apply a 100m **buffer** around all **stands** of modelled **Ash old growth forest** that are depicted in the **Department**'s corporate spatial datasetMOG2009.shp, and confirmed during field assessment by the **managing authority** or the **Department** to be **Ash** type **forest**.
- 4.3.5 White Cypress Pine, Buloke, Grey Box, Yellow Box and Grey Box vegetation communities
- 4.3.5.1 In the Mid Murray **FMA**, exclude **timber harvesting operations** within any White Cypress Pine (*Callitris glaucophylla*), Buloke, Grey Box (*Eucalyptus melliodora*), Yellow Box (*Eucalyptus melliodora*) and Grey Box vegetation communities.

4.3.6 Snow Gum and Swamp Gum

4.3.6.1 For vegetation communities dominated by Snow Gum (*Eucalyptus pauciflora*) on Mt Cole and Swamp Gum (*Eucalyptus ovata*) near Enfield in the Midlands **FMA** that aren't already classified as **SPZ**, apply a **protection area** prior to commencement of the **timber harvesting operation** consistent with management actions listed in **Table 15 Detection based rules for Heathland, Snow Gum, Swamp Gum and Heath vegetation communities**.

4.3.7 Rainforest canopy species

4.3.7.1 **Rainforest canopy** species are defined as shade tolerant tree and vine species which are able to regenerate below an undisturbed **canopy**, or in small **canopy** gaps resulting from

locally recurring minor disturbances, such as isolated windthrow or lightning strike, which are part of the **rainforest** ecosystem. Such species are not dependent on fire for their **regeneration**.

- 4.3.7.2 In East Gippsland, Warm Temperate **Rainforest canopy** species are: Lilly Pilly (*Acmena smithii*), Kanuka (*Tristaniopsis laurina*), Sweet Pittosporum (*Pittosporum undulatum*), Blackwood (*Acacia melanoxylon*), Blue Olive-berry (*Elaeocarpus reticulates*), Muttonwood (*Myrsine howittiana*), Jungle Grape (*Cissus hypoglauca*), Boobialla (*Myoporum insulare*) and Yellow-wood (*Acronychia Ablongifolia*).
- 4.3.7.3 In East Gippsland, Cool Temperate **Rainforest canopy** species are: Southern Sassafras (*Atherosperma moschatum*), Black Olive-berry (*Elaeocarpus holopetalus*), Blackwood (*Acacia melanoxylon*), Gippsland Waratah (*Telopea oreades*), Privet Mock-olive (*Notelaea ligustrina*), Banyalla (*Pittosporum bicolor*), Goonmirk Rocks Errinundra Plum Pine (*Podocarpus sp. aff. Lawrence*) and Errinundra Pepper (*Tasmannia xerophila subsp. robusta*).
- 4.3.7.4 In the **Central Highlands FMAs** and Central Gippsland **FMA**, Cool Temperate **Rainforest canopy** species are: Myrtle Beech (*Nothofagus cunninghamii*), Southern Sassafras (*Atherosperma moschatum*), Blackwood (*Acacia melanoxylon*), Banyalla (*Pittosporum bicolor*), Mountain Pepper (*Tasmannia lanceolata*) and Mountain Tea-tree (*Leptospermum grandifolium*).
- 4.3.7.5 In the Central Gippsland **FMA**, Warm Temperate **Rainforest canopy** species are: Sweet Pittosporum (*Pittosporum undulatum*), Muttonwood (*Myrsine howittiana*) and Blackwood (*Acacia melanoxylon*).
- 4.3.7.6 In the Otway's, Cool Temperate **Rainforest canopy** species are: Myrtle Beech (*Nothofagus cunninghamii*) and Blackwood (*Acacia melanoxylon*).
- 4.3.7.7 In all areas, **forest stands** that are dominated by single (contain over 50%), non-diverse **stands** of the following species and which contain few other **rainforest** characteristics, are not considered **rainforest**: Blackwood (*Acacia melanoxylon*), Mountain Teatree (*Leptospermum grandifolium*), Sweet Pittosporum (*Pittosporum undulatum*), Errinundra Pepper (*Tasmannia xerophila subsp. robusta*) or Mountain Pepper (*Tasmannia lanceolata*).
- 4.3.7.8 Eucalypts are not classified as broad-leaved **rainforest** species and therefore Eucalypts do not contribute to the projected foliage cover when identifying **stands** of **rainforest**.
- 4.3.8 Rainforest field recognition and delineation
- 4.3.8.1 **Rainforest stands** are identified in the field as **forest** of at least 0.1 hectares where the projected foliage cover of the broadleaved tree **canopy** is greater than 70 % and is contributed by 1 or more of the **canopy** tree species listed above in sections 4.3.7.2 to 4.3.7.6 and where section 4.3.7.7 is not met.
- 4.3.8.2 Projected foliage cover is the proportion of ground covered by the vertical projection of foliage and branches from **rainforest canopy** trees. Any potential contribution of **understorey** species such as tree ferns is not counted toward assessment of projected foliage cover.
- 4.3.8.3 Linear **stands** are defined as **stands** of **rainforest** which are elongated and which are between 20m and 40m wide. Linear **stands** of **rainforest** usually occur along **drainage lines** or small streams. Linear **stands** may be "overshadowed" by eucalypts from the adjoining eucalypt **forest**.
- 4.3.8.4 The minimum area for recognition of a **rainforest stand** is 0.1 ha and the minimum narrowest width is 20m. (i.e. 20m by 50m).
- 4.3.8.5 Special care is required when assessing the boundary between identified **rainforest stands** and adjacent wet sclerophyll **forest stands** where disturbance such as fire has temporarily removed the **rainforest canopy** or has created temporary **canopy** gaps within the **stand** of **rainforest**. In cases where the **canopy** disturbance is less than ten years old and further

guidance as to the boundary between identified **rainforest** and adjacent wet sclerophyll **forest stands** is required, the 'differential species approach' is to be used (Differential species keys for the delineation of **rainforest** boundaries can provide reference photos)

- 4.3.8.6 Where the **rainforest canopy** is absent and there is little or no evidence of the **regeneration** of a **rainforest canopy** after 10 years following disturbance the 'differential species approach' should not be used to identify **rainforest** and the **stand** should no longer be considered to be **rainforest**.
- 4.3.8.7 Where the 'differential species approach' is utilised, the **rainforest** boundary is the point where the number of **rainforest** species equals the number of eucalypt **forest** species i.e. the line along which the **floristic** signals are of equal strength. This approach would be used where the **rainforest canopy** tree cover reduces gradually from 70% projected foliage cover. (gradual transition is a transition from 70% **rainforest** species projected foliage cover to 70%non-**rainforest** species projected foliage cover over a distance greater than approximately 10 meters)
- 4.3.9 Rainforest protection measures
- 4.3.9.1 Protect all **rainforest stands** from **timber harvesting operations** as follows:
 - a) Exclude non linear **stands** that are 0.1ha or more in size but less than 0.4ha from **timber harvesting operations**. These **stands** do not require a **buffer**.
 - b) Exclude linear **stands** that are at least 0.1ha but are less than 0.2ha from **timber harvesting operations**. These **stands** do not require a **buffer**.
 - c) Exclude linear **stands** that are at least 0.2ha but are less than 0.4ha from **timber harvesting operations**. Protect these **stands** with a 20m **buffer**.
 - d) Exclude all **rainforest stands** (including linear **stands**) equal to or exceeding 0.4ha from **timber harvesting operations**. Protect these **stands** with a 40m **buffer** except for **rainforest stands** in the **Central Highlands FMAs** and the **Gippsland FMAs** where 4.3.9.2 below must be complied with.
 - e) Distribute slash away from retained rainforest stands or buffers.
- 4.3.9.2 In areas categorised as being of National, State or Regional significance in the **Sites of Significance for Rainforest** spatial layer where evidence of **rainforest** greater in size than 0.4ha is found in the field and it isn't already classified as **SPZ**, apply a **protection area** prior to commencement of the **timber harvesting operation** consistent with management actions listed in **Table 16 Buffer widths for Rainforest Sites of Significance by category and priority**.

4.4 Pests, weeds and diseases

4.4.1 Diseases

- 4.4.1.1 Minimise the risk of introduction or movement of Cinnamon Fungus (*Phytophthora cinnamomi*) and Root Rot (*Armillaria*) from a **forest disease control area** into other areas, by:
 - a) washing machinery before moving from a forest disease control area into other areas;
 - b) restricting activities where the movement of soil or gravel is likely to cross from a **forest disease control area** into other areas:
 - c) minimising the relocation or movement of gravel or soil during **road** and track construction or maintenance works, or logging operations within a **forest disease control area**;
 - d) restricting or controlling drainage water run-off from roads and tracks to minimise run-off from a forest disease control area into other areas;
 - e) testing gravel from **forest disease control areas** and using only uncontaminated gravel in other areas; and
 - f) cleaning and disinfecting vehicles, machinery, tools and equipment used within a forest disease control area.
- 4.4.1.2 Minimise the spread of Myrtle Wilt (*Chalara australis*) when operating in a **forest disease control area** by:
 - a) protecting individual Myrtle Beech (Nothofagus cunninghamii) trees;
 - b) sterilising equipment with anti-fungal agent or warm water and soap prior to moving into a new area;
 - c) pruning Myrtle Beech (*Nothofagus cunninghamii*) that are subject to ongoing damage by vehicles; and
 - d) immediately treating wounds on Myrtle Beech (*Nothofagus cunninghamii*) (including those left by pruning) with a commercial, waterproof wound sealant.

4.4.2 Weeds

- 4.4.2.1 Conduct a pre-harvest assessment to determine the type and extent of weeds on the **coupe** and on associated **coupe access roads**.
- 4.4.2.2 Conduct a post-harvest assessment to determine the type and extent of weeds on the **coupe** and associated **coupe access roads** in the first spring after completion of **site preparation** and establishment and during the **stocking** survey.
- 4.4.2.3 Where the assessments identify the **timber harvesting operation** has introduced or exacerbated weeds in a **coupe**, prepare a weed management plan and implement a weed control program.
- 4.4.2.4 Record any areas to be treated on a map in the **Forest Coupe Plan** and mark in the field as necessary prior to treatment.

5. Important values

5.1 Heritage, historic places and Aboriginal cultural heritage

- 5.1.1.1 During planning for **timber harvesting operations**, have regard to obligations under relevant heritage legislation including the:
 - i. Heritage Act 2017,
 - ii. Aboriginal Heritage Act 2006.
- 5.1.1.2 Ensure relevant databases are consulted prior to the commencement of **timber harvesting operations** including the:
 - i. Victorian Heritage Register,
 - ii. Victorian Heritage Inventory,
 - iii. Aboriginal Cultural Heritage Register and Information System (ACHRIS) public map,
 - iv. Forest Management Zoning Scheme.
- 5.1.1.3 Where a **historic place** or **Aboriginal cultural heritage** is present and represented within the **Forest Management Zoning Scheme**, comply with zoning and management actions. Where a **historic place** or **Aboriginal cultural heritage** is present and not represented within the **Forest Management Zoning Scheme**, relevant heritage laws may still apply.
- 5.1.1.4 Where there is evidence of the presence of a **historic place** or **Aboriginal cultural heritage** that is not or may not be registered under relevant heritage legislation, ensure the evidence of the **historic place** or **Aboriginal cultural heritage** is reported to the **Secretary** and to any person or body responsible for the relevant legislation, and the **historic place** or **Aboriginal cultural heritage** is protected from **timber harvesting operations** (subject to obtaining any relevant permission under the applicable heritage legislation).

5.2 Landscape Sensitivity Areas

- 5.2.1 Central Highlands FM As
- 5.2.1.1 Retain all **mature** trees within 20m of the Monda Track.
- 5.2.1.2 Apply a 50m **buffer** either side of La La Falls walking track.
- 5.2.1.3 Apply a 50 m **buffer** either side of Island Creek walking track and a 100m **buffer** around the Ada Tree.
- 5.2.1.4 Apply an 80m **buffer** upslope and 20m **buffer** downslope of aqueducts in the Historic and Cultural Features Reserves in Rubicon State Forest.

Foreground Seen Area Management (0 – 500 m)

5.2.1.5 In the foreground, within 500m seen from the 'Places' listed as 'Landscape Features', 'Scenic Drives' and 'Designated Lookouts' in **Table 17 Landscape management prescriptions**,

manage **timber harvesting operations** to ensure landscape alterations are temporary, subtle and not evident to the casual observer.

5.2.1.6 In the foreground, within 500m seen from the 'Places' listed as 'Scenic Drives' and 'Designated Lookouts' in **Table 17 Landscape management prescriptions**, use a minimum 20m vegetation **buffer** to screen **timber harvesting operation**s (except **selective harvesting** operations) and new **road** alignments from view.

Middleground seen area management (500m – 6.5km)

- 5.2.1.7 In the middleground, between 500m and 6.5km, seen from the 'Places' listed as 'Landscape Features' in **Table 17 Landscape management prescriptions**:
 - (a) manage **timber harvesting operations** to ensure landscape alterations are only subtly apparent within 5 years of the operation; and
 - (b) shape, position and time **timber harvesting operations** and new **roads** to minimise their visual impact.

5.2.2 East Gippsland FMA

5.2.2.1 Within the 'Places' listed as 'Landscape Features' described in **Table 17 Landscape** management prescriptions, limit the area of **coupes** which are highly visible from Gelantipy Road to 20ha and leave at least 3 years between harvesting of adjacent **coupes**.

Foreground seen area management (0 – 500 m)

- 5.2.2.2 In the foreground, within 500m seen from the 'Places' listed as 'Landscape Features', 'Scenic Drives' and 'Designated Lookouts' listed in **Table 17 Landscape management** prescriptions, manage **timber harvesting operations** to ensure landscape alterations are temporary, subtle and not evident to the casual observer.
- 5.2.2.3 In the foreground, within 500m seen from the 'Places' listed as 'Scenic Drives' and 'Designated Lookouts' in Table 17 Landscape management prescriptions, use a minimum 20m vegetation buffer to screen timber harvesting operations (except selective harvesting operations) and new road alignments from view.

Middleground seen area management (500m – 6.5km)

- 5.2.2.4 In the middle ground, between 500m and 6.5km, seen from the 'Places' listed as 'Scenic Drives' and 'Designated Lookouts' in **Table 17 Landscape management prescriptions**:
 - a) manage timber harvesting operations to ensure landscape alterations are only subtly apparent within 2 years of the operation; and
 - b) shape, position and time **timber harvesting operation**s and new **roads** to minimise their visual impact.

Background ridgeline management

5.2.2.5 Design **timber harvesting operations** so that **clearfall** and skyline 'notches' are not apparent from the 'Places' listed as 'Scenic Drives' and 'Designated Lookouts' listed in **Table 17 Landscape management prescriptions**.

5.2.3 Gippsland FMAs

5.2.3.1 Plan and implement **timber harvesting operations** in areas zoned as **SMZ** for landscape values in a manner that ensures the scenic value is maintained.

5.2.3.2 In the middle ground seen from the 'Places' listed as 'Landscape Features' listed in **Table 17 Landscape management prescriptions**, design **timber harvesting operations** and new **roads** to minimise their visual impact.

5.2.4 Midlands FMA

Foreground seen area management (0 – 500m)

- 5.2.4.1 In the foreground, within 500m seen from the 'Places' listed as 'Scenic Drives' and 'Designated Lookouts' listed in **Table 17 Landscape management prescriptions** manage **timber harvesting operations** to ensure landscape alterations are temporary, subtle and not evident to the casual observer.
- In the foreground, within 500m seen from the 'Places' listed as 'Scenic Drives' and 'Designated Lookouts' in **Table 17 Landscape management prescriptions**, use a minimum 20m vegetation **buffer** to screen **timber harvesting operations** (except **selective harvesting** operations) and new **road** alignments from view.

Middleground seen area management (500m – 6.5km)

- 5.2.4.3 In the middle ground, between 500m and 6.5km, seen from the 'Places' listed as 'Scenic Drives' and 'Designated Lookouts' in **Table 17 Landscape management prescriptions**:
 - a) manage **timber harvesting operations** to ensure landscape alterations are only subtly apparent within 2 years of the operation; and
 - b) shape, position and time **timber harvesting operations** and new **roads** to minimise their visual impact.

5.2.5 Mid Murray FMA

- 5.2.5.1 Within the landscape management **SMZ**s, manage **timber harvesting operations** as follows:
 - a) retain large old trees close to the roads (subject to public safety);
 - b) during peak periods of visitor use, conduct only those **timber harvesting operations** and silvicultural works (like **thinning**) which maintain visual amenity; and
 - c) ensure harvesting debris is removed from the corridor.
- 5.2.5.2 Site log **landings** and sleeper cutting ducks at least 60m from **roads** subject to landscape management **SPZ**s.

5.2.6 North East FMAs

- 5.2.6.1 For Indicative National Estate natural landscapes, the following management directions apply:
 - a) At Mt. Cudgewa/Rogers Creek, design coupes to minimise impact on view from Henlow (Tallangatta Valley);
 - b) At Lower Howqua North/Red Hill, design **coupes** to minimise impact on view from Mt. Buller:
 - At Mt. View (Snake-Edwards Divide), design coupes to minimise impact on view from Mt. Skene Road.
- 5.2.6.2 For the Indicative National Estate values at Eskdale Spur and Mt Emu, design **coupe** to minimise impact on view from Kiewa Valley Highway.
- 5.2.6.3 For the Indicative National Estate value (natural, aesthetic) along Omeo Highway and at Mt. Wills Historic Area:

- a) Plan timber harvesting operations between Bogong Saddle and Glen Wills on the Omeo Highway to ensure no more than 10% of the length of roadside should be less than 3 years old; and
- b) Harvesting adjacent to or immediately opposite a recently harvested **coupe** may only proceed when the **regeneration** of the initial harvested area is at least 3 years old.
- 5.2.6.4 Plan and implement **timber harvesting operations** and new **road construction** in existing landscape **SMZ** in a manner that ensures the scenic value is maintained.
- 5.2.6.5 In the middle ground seen from the 'Places' listed as 'Landscape Features' in **Table 17 Landscape management prescriptions**, design **timber harvesting operations** and new **roads** to minimise their visual impact.

5.3 Apiary

5.3.1 Bendigo FMA

5.3.1.1 When planning harvesting in patches of summer-flowering Red Ironbark **stands**, ensure that potential short-term flowering reductions resulting from harvesting are not concentrated in time or location.

5.3.2 Gippsland Forest Apiary Plan Area

- 5.3.2.1 Within a 1km radius of Apiary Zone 1 sites (sites of high importance to the apiary industry), manage **timber harvesting operations** to:
 - a) maximise the proportion of older age **stands**;
 - b) retain at least 66% of **State forest** at more than 40 years of age;
 - retain at least 66% of Yellow Stringybark (Eucalyptus muelleriana) and Red Box (Eucalyptus polyanthemos) stands in State forest in the mature and/or over-mature age class; and
 - d) retain all individual Blue Box (Eucalyptus baueriana) trees.

5.4 Recreation

- 5.4.1.1 For developed recreation facilities not included in the **SMZ** or **SPZ** components of the **FMZS**, exclude **timber harvesting operations** within 40m. Recreation facilities are considered to be developed if they include fixed structures, campsites, and/or signage that is maintained by the **land manager** or an authorised operator.
- 5.4.1.2 For the recreation sites listed in **Table 18 Recreation sites**, document and **manage timber** harvesting operations to maintain the value of the place.
- 5.4.1.3 Within the Australian Alps Walking Track (AAWT) **SPZ**:
 - (a) construct and maintain **roads** that cross the AAWT in a manner that considers the needs of walkers using the track; and
 - (b) consider the use of alternative extraction routes during harvesting operations to protect the AAWT.

5.5 Research

- 5.5.1.1 In the **Gippsland FMAs**, Midlands **FMA** and **North East FMAs** refer to **Table 19 Research** sites for prescriptions applicable to research locations and projects.
- 5.5.1.2 In Mid Murray **FMA**, exclude **timber harvesting operations** that are incompatible with the objectives for **sanctioned** research projects, until such time as the areas are no longer required.
- 5.5.1.3 In the **Central Highlands FMA**, exclude **timber harvesting operations** within 50m of registered ANU Monitoring sites at which long-term ecological monitoring is being conducted.

5.6 Fuel Hazard ratings in Fire Management Zones

- 5.6.1.1 The scheduling of **timber harvesting operations** and **planned burning** should occur to ensure that areas are **clearfall** or **seed tree** harvested less than 12 months prior to planned burning.
- 5.6.1.2 Failure to comply with clauses 5.6.1.3 through to 5.6.1.6 may result in the **Department** undertaking planned burning in the **coupe**. Where this occurs, the **managing authority** or **harvesting entity** is responsible for ensuring **regeneration** standards are met post planned burning.
- 5.6.1.3 The time frame for ensuring that the overall **fuel hazard ratings** are met commences from **coupe** completion (i.e. not temporary closure). A maximum of 2 years from **coupe** commencement is allowed if temporary clearances are applied to a **coupe**.
- 5.6.1.4 The following requirements apply to **APZ Land**:
 - a) The overall fuel hazard rating for a harvested coupe must be less than or equal to moderate:
 - i. during harvesting if harvesting is conducted during the fire season; and
 - ii. within 12 months of harvesting.
 - b) Clearfall and seed tree harvesting is not permitted in more than 1_ha of the gross coupe area.
 - c) Harvesting using **thinning** or **selective harvesting** must not be undertaken if it would cause the rolling five year average APZ harvest percentage within a **fuel hazard management unit** to exceed 5 per cent.
 - d) For the purposes of paragraph (c), the rolling five year average APZ harvest percentage is to be determined in the 2022/23 financial year and each subsequent financial year, for each **fuel hazard management unit**, using the following method:
 - i. Calculate the annual APZ harvest percentage in a **fuel hazard management unit** for the financial year using the following formula:

$$A = \left(\frac{H}{Z}\right) \times 100$$

Where -

A is the annual APZ harvest percentage;

H is the total combined area, measured in hectares, of **APZ Land** in the **fuel hazard management unit** that was zoned either **GMZ** or **SMZ** as of 30 June of the preceding financial year, which has been harvested using **thinning** or **selective harvesting** in the financial year;

Z is the total combined area, measured in hectares, of **APZ Land** in the **fuel hazard management unit** that was zoned either **GMZ** or **SMZ**, as at 30 June of the preceding financial year.

ii. Add the annual APZ harvest percentage for the financial year to the final annual APZ harvest percentage of each of the immediately preceding four financial years, and divide that total by 5.

Note: the annual APZ harvest percentage formula applies continuously throughout a financial year (including to be calculated at the conclusion of the financial year, taking into account harvesting up to and including 30 June of that year, which will be the final annual APZ harvest percentage for that year).

- e) For the purposes of paragraph (d)(ii), the final annual APZ harvest percentage for the 2018/19, 2019/20, 2020/21 and 2021/22 financial years must be calculated:
 - i. as soon as practicable after 1 July 2022 (and in any event no later than 1 September 2022);
 - ii. using the method specified in paragraph (d) (as modified by paragraphs (iii) and (iv) below);
 - iii. on the assumption that **fuel hazard management units** were in place and that had the same boundaries as each **fuel hazard management unit** has as at 1 July 2022.
 - iv. on the assumption that, within a **fuel hazard management unit**, the boundaries of the **Asset Protection Zones**, **GMZ** and **SMZ** were the same as those that exist as at 1 July 2022.

5.6.1.5 The following requirements apply to **BMZ Land**:

- a) The **overall fuel hazard** rating for a **clearfall** or **seed tree harvested coupe** must be returned to a rating of less than or equal to high within 12 months of harvesting.
- b) Harvesting using **clearfell** or **seed tree harvesting** must not be undertaken if it would cause the rolling five year average BMZ clearfell harvest percentage within a **fuel hazard management unit** to exceed 1 per cent.
- c) For the purposes of paragraph (c), the rolling five year average BMZ clearfell harvest percentage is to be determined in the 2022/23 financial year and each subsequent financial year, for each fuel hazard management unit, using the following method:
 - i. Calculate the annual BMZ clearfell harvest percentage in a **fuel hazard management unit** for the financial year using the following formula:

$$BC = \left(\frac{H}{Z}\right) \times 100$$

Where -

BC is the annual BMZ clearfell harvest percentage;

H is the total combined area, measured in hectares, of **BMZ Land** in the **fuel hazard management unit** that was zoned either **GMZ** or **SMZ** as of 30 June of the preceding financial year, which has been harvested using clearfell or seed tree harvesting in the financial year;

Z is the total combined area, measured in hectares, of **BMZ** Land in the fuel hazard management unit that was zoned either **GMZ** or **SMZ**, as at 30 June of the preceding financial year.

ii. Add the annual BMZ clearfell harvest percentage for the financial year to the final annual BMZ clearfell harvest percentage of each of the immediately preceding four financial years, and divide that total by 5.

Note: the annual BMZ clearfell harvest percentage formula applies continuously throughout a financial year (including to be calculated at the conclusion of the financial year, taking into account harvesting up to and including 30 June of that year, which will be the final annual BLMZ clearfell harvest percentage for that year).

- d) For the purposes of paragraph (c)(ii), the final annual BMZ clearfell harvest percentage for the 2018/19, 2019/20, 2020/21 and 2021/22 financial years must be calculated:
 - i. as soon as practicable after 1 July 2022 (and in any event no later than 1 September 2022).
 - ii. using the method specified in paragraph (c) (as modified by paragraphs (iii) and (iv) below);
 - iii. on the assumption that **fuel hazard management units** were in place and had the same boundaries as each **fuel hazard management unit** has as at 1 July 2022.
 - iv. on the assumption that, within a **fuel hazard management unit**, the boundaries of the Bushfire Moderation Zones, **GMZ** and **SMZ** were the same as those that exist as at 1 July 2022.
- e) Harvesting using **thinning** or **selective harvesting** must not be undertaken if it would cause the rolling five year average BMZ thinning harvest percentage within a **fuel hazard management unit** to exceed 4 per cent.
- f) For the purposes of paragraph (e), the rolling five year average BMZ thinning harvest percentage is to be determined in the 2022/23 financial year and each subsequent financial year, for each **fuel hazard management unit**, using the following method:
 - i. Calculate the annual BMZ thinning harvest percentage in a **fuel hazard management unit** for the financial year using the following formula:

$$BT = \left(\frac{H}{Z}\right) \times 100$$

Where -

BT is the annual BMZ thinning harvest percentage;

H is the total combined area, measured in hectares, of BMZ Land in the fuel hazard management unit that was zoned either GMZ or SMZ as of 30 June of the preceding financial year, which has been harvested using thinning or selective harvesting in the financial year:

Z is the total combined area, measured in hectares, of **BMZ** Land in the fuel hazard management unit that was zoned either **GMZ** or **SMZ**, as at 30 June of the preceding financial year.

ii. Add the annual BMZ thinning harvest percentage for the financial year to the final annual BMZ thinning harvest percentage of each of the immediately preceding four financial years, and divide that total by 5.

Note: the annual BMZ thinning harvest percentage formula applies continuously throughout a financial year (including to be calculated at the conclusion of the financial year, taking into account harvesting up to and including 30 June of that year, which will be the final annual BMZ thinning harvest percentage for that year).

- g) For the purposes of paragraph (f)(ii), the final annual BMZ thinning harvest percentage for the 2018/19, 2019/20, 2020/21 and 2021/22 financial years must be calculated:
 - i. as soon as practicable after 1 July 2022 (and in any event no later than 1 September 2022).
 - ii. using the method specified in paragraph (f) (as modified by paragraphs (iii) and (iv) below);
 - iii. on the assumption that **fuel hazard management units** were in place and had the same boundaries as each **fuel hazard management unit** has as at 1 July 2022.
 - iv. on the assumption that, within a **fuel hazard management unit**, the boundaries of the Bushfire Moderation Zones, **GMZ** and **SMZ** were the same as those that exist as at 1 July 2022.
- h) **Thinned** or **selectively harvested coupes** may be burnt by the **Department** as part of future planned burns. These burns will be conducted at least 18 months post harvest.
- 5.6.1.6 If any part of a **coupe** is within 1.5km of private property, and any part of that **coupe** is also within a Landscape Management Zone as described in the *Code of Practice for Bushfire Management on Public Land 2012*, the overall **fuel hazard rating** for that **coupe** if harvested (unless harvested using **selective harvesting** or **thinning**) must be less than or equal to high within 12 months of harvesting.

6. Roading

6.1	Road planning
6.1.1.1	Plan new roads and significant road improvement operations to minimise construction through management areas , SMZ , protection areas , SPZ , wet, unstable areas, and slopes greater than 30 degrees.
6.1.1.2	Application may be made to the Minister or delegate to obtain approvals for road construction (including coupe access and in-coupe roads) or significant road improvement operations proposed to be roading activities conducted in protection areas or SPZ in accordance with Section 1.4.
6.1.1.3	Seek engineering advice for road alignments traversing cross slopes of 30 degrees or greater or 25 degrees and greater in areas of high soil erodability.
6.1.1.4	Identify the intended class of a new road or road upgrade in accordance with the appropriate service function description in Table 20 Road classification system
6.1.1.5	Design new roads and road upgrades to conform to the geometric design standards in Table 21 Guidelines to the main geometric design standards – unsealed roads for the intended road class.
6.1.2	Clearing Widths
6.1.2.1	Minimum clearing widths for roads are specified in Table 22 Minimum clearing widths (m) required for typical road construction
6.1.2.2	When planning clearing widths for road intersections on permanent roads include the minimum formation width plus any additional width required for the construction of batters.
6.1.2.3	Where a slashed verge is necessary, plan clearing widths for permanent roads sufficiently wide to enable efficient control of unwanted regrowth .
6.1.2.4	Limit clearing widths to those specified in Table 22 Minimum clearing widths (m) required for typical road construction plus any additional width required to construct batters.

6.2	Road Construction
6.2.1.1	Undertake road construction when rainfall and soil conditions minimise the risk of erosion and impact on water quality, and when soil moisture is adequate to achieve compaction and stabilisation of the sub-grade.
6.2.1.2	Clear road alignments prior to road formation.
6.2.1.3	For construction of permanent roads , remove all stumps, logs and other debris from within the formed width of the road site.
6.2.1.4	Where road construction requires the removal of topsoil in large quantities maintain the topsoil in a stockpile where practicable, clear of logging debris for use in batter stabilisation, snig track rehabilitation or other coupe infrastructure rehabilitation.
6.2.1.5	Create table drains by extending the road when it is formed, and not by subsequent excavation.
6.2.1.6	Limit earthworks to the least possible to achieve the road design specification.
6.2.1.7	At the completion of permanent road construction operations, notify the Department of the location of the road (so that the road may be included on the Department Road Register).
6.2.2	Fill Batter Construction
6.2.2.1	Minimise fill batters from covering the base of live retained trees .
6.2.2.2	Only use clean and weed free mulch in batter rehabilitation works.
6.2.2.3	Stabilise fill batters using mechanical consolidation where practical, to manage any soil movement away from the fill area.
6.2.3	Surfacing
6.2.3.1	Consolidate sub-grades before placing pavement material.
6.2.3.2	Consolidate and level the base course material prior to placing the wearing course material.
6.2.3.3	On permanent roads use surfacing materials appropriate to passenger vehicles and timber harvesting operations .
6.2.4	Road Drainage
6.2.4.1	The maximum distance between drainage structures for road grade and soil erosion hazard is specified in Table 23 Maximum distance between drainage structures .
6.2.4.2	Construct cross-drains at an angle sufficient to discharge any water from the surface of the road .
6.2.4.3	On soils of high erosion hazard, use temporary sediment traps to prevent erosion during road construction .
6.2.4.4	Appropriate discharge areas for drainage include:
	a) a strip of undisturbed vegetation at least 20 m wide;
	b) a rock spill; or
	c) some other structure that dissipates the velocity of drainage flows.

- 6.2.4.5 Place drainage structures at least 20 m from permanent or **temporary streams**, to allow discharge onto undisturbed vegetation and to maximise the flow distance between the drainage outlet and the **waterway**.
- 6.2.4.6 Where it is not practical to comply with clause 6.2.4.5, and a drainage structure must be located within 20 m of a permanent or **temporary stream**:
 - a) use crown or cross fall techniques to drain roads into undisturbed vegetation; or
 - b) pass drainage through an appropriate sediment control structure such as a sediment pond or silt trap before entering a permanent or **temporary stream**.
- 6.2.4.7 Construct table drains to:
 - a) allow water to flow, without ponding;
 - b) include run-offs of sufficient length to allow the table drain and run-offs to be cleaned;
 - c) be supported by rock or otherwise stabilised in soils of a high erosion hazard; and
 - d) have silt traps constructed at the end if discharging directly into a stream or **wetland buffer**.
- 6.2.5 Culverts
- 6.2.5.1 Culverts used in **permanent roads** are a minimum of 375 mm in **diameter**.
- 6.2.5.2 Culverts used in **temporary roads** are a minimum of 300mm in **diameter**.
- 6.2.5.3 All culverts are designed to withstand a 1 in 10 year rainfall event.
- 6.2.5.4 Construct culverts in catchment areas exceeding 100ha in accordance with engineering advice.
- 6.2.5.5 On **drainage lines**, stream and river crossings or soils of High Erosion Hazard place sandbags, **timber**, concrete, or rock at the head of the culvert and at the point of discharge to hold the culvert in place and protect it from erosion.
- 6.2.5.6 Include a **road** sump for all culverts on Class 5C and higher **roads**.
- 6.2.5.7 If constructed of concrete, have a minimum cover of 600 mm as measured from the **road** surface to the top of the pipe and a maximum cover as specified in the Installation of Steel-Reinforced Concrete Drainage Pipelines, Concrete Pipe Association of Australasia.
- 6.2.5.8 If constructed of a material other than concrete, have a minimum cover over the pipe as recommended in the manufacturer's specifications.
- 6.2.5.9 On **permanent streams**, include a fish ladder if the **diameter** of the culvert is greater than 750 mm.
- 6.2.5.10 Protect any fill face upstream or downstream of a culvert in a way that prevents erosion.
- 6.2.5.11 Ensure culverts do not project above the bed of a **waterway** in a way which may prevent the passage of aquatic **fauna**.
- 6.2.5.12 Where culvert construction diverts water from its natural course, return water to its natural course over a flume, rock spill, or other hard surface.

6.2.6	Bridges
6.2.6.1	Design bridges in accordance with the Department's Bridge and Major Culvert Policy.
6.2.6.2	Design bridges to prevent constriction of any clearly defined channel.
6.2.6.3	Source earth borrow from outside waterway buffers.
6.2.6.4	Ensure that excavations, sills, abutments, stringers, and girders are made or placed above the high watermark of the stream, wetland or drainage line .
6.2.6.5	Protect bridges from erosion by use of natural groundcover, a retaining wall, a bulkhead, or a rock surface.
6.2.6.6	Only construct temporary crossings to carry machinery during bridge construction if the bed of the stream, wetland or drainage line is capable of bearing the weight of that machinery without being damaged.
6.2.6.7	Temporary crossings to carry machinery during bridge construction:
	a) include a corduroy crossing of logs;
	b) are adequately drained, along with any access tracks, when construction is complete; and
	c) are removed and rehabilitated on completion of works.
6.2.7	Fords
6.2.7.1	The base and entry points of fords are constructed of rock, concrete, heavy timber or other erosion-resistant material.
6.2.7.2	Fords are only as wide as the crossing place will allow.
6.2.7.3	Ensure fords do not project above the bed of a waterway in a way which may prevent the passage of aquatic fauna .
6.3	Road Maintenance
6.3.1.1	Breach at regular intervals any soil windrow erected on the outside of a road .
6.3.1.2	Maintain drainage structures to minimise the risk that accumulation of debris and run-off could results in damage to drainage structures.
6.4	Road Rehabilitation
6.4.1.1	Close temporary roads (including removal of all bridges, crossings and culverts on streams or drainage lines) as soon as possible after harvesting and/or regeneration is complete in all coupes that use the road .
6.4.1.2	Drain the approach to any bridge, culvert or log fill crossing that has been removed to restrict soil movement into a stream or waterway .
6.4.1.3	Use an effective barrier to close to all vehicles temporary roads that will not be used to access

a **coupe** for a period of 12 months or more.

7. Timber Harvesting

7.1 Coupe Management

- 7.1.1 Coupe Boundary
- 7.1.1.1 The **coupe** boundary must be able to be identified in the field and must reflect the mapped **coupe** boundary on the **sanctioned Forest Coupe Plan**.
- 7.1.1.2 Where a **coupe** boundary shown in a **Forest Coupe Plan** is mapped to a geographic feature and the actual location of that geographic feature does not match the mapped location, the **coupe** boundary may be moved a maximum of 50m from the mapped boundary to align with the actual location of the intended boundary feature. If the distance exceeds 50m then the **Forest Coupe Plan** Map must be amended to reflect the actual **coupe** boundary.
- 7.1.1.3 Where the **coupe** boundary abuts non-**State forest** land tenures, the **managing authority** must consult with the adjoining **land managers** / **land owners** and agree on the location of the mapped **coupe** boundary in the field.
- 7.1.2 Areas where timber harvesting operations are to be excluded
- 7.1.2.1 **Timber harvesting operations** must be excluded from:
 - a) SPZ or protection areas;
 - b) exclusion areas within SMZ;
 - c) exclusion areas; and
 - d) within 10m of vertical or near vertical sided gullies with a depth of half a metre or more that are actively eroding (or within 20m where slope exceeds 20 degrees) in the Bendigo **FMA**.
- 7.1.2.2 Where an **exclusion area** or **protection area** is based on an incorrectly mapped geographic feature or value, the **exclusion area** or **protection area** should be centred on the actual location of the feature or value in the field not the previously mapped location.
- 7.1.2.3 **Exclusion areas, protection areas** and **SPZs** must be protected from damage during rough heaping or windrowing operations.
- 7.1.2.4 Application may be made to the **Minister** or **delegate** in accordance with Section 1.4 to conduct the following **timber harvesting operations** in **Special Protection Zones** and **protection areas**:
 - a) regeneration or rehabilitation works in areas of harvested native forest; or
 - b) use of existing **coupe infrastructure** (and necessary incidental works) in areas of harvested **native forest**.

7.1.3 Operations in buffers

- 7.1.3.1 Trees can be harvested within **buffer** areas if **sanctioned** for safety purposes.
- 7.1.3.2 Machinery is to be excluded from **buffers** except where involved in the **sanctioned** construction of a **coupe access road**, **in-coupe road**, or stream crossing; or when using an established stream crossing or **existing road** for the purpose of **road maintenance** works, **snigging**, or **haulage**.
- 7.1.3.3 Except where constructing a **sanctioned** stream crossing, minimise circumstances where fill and harvesting debris may enter **buffers**.

7.1.4 Operations in filters

- 7.1.4.1 Machinery is to be excluded from **filter strips** except where involved in the **sanctioned** construction of a **coupe access road**, **in-coupe road**, stream crossing or when using an established stream crossing or **existing road** for the purpose of **road maintenance** works, **snigging** or **haulage**.
- 7.1.4.2 Except where constructing a **sanctioned** stream crossing, minimise circumstances where fill and harvesting debris may enter **filter strips**.
- 7.1.4.3 **Filter strips** must be protected from damage during rough heaping or windrowing operations.

7.2 Coupe Infrastructure

- 7.2.1 Snig Track and Landing Construction
- 7.2.1.1 Crossing standards and procedures for **roads** also apply to **snig track** crossings.
- 7.2.1.2 Avoid placing bark on uncorded **snig tracks** (this does not apply to **thinning outrows**).
- 7.2.1.3 Where **cording** is to be used, it should not be placed on **snig tracks** if machinery caused soil damage already exists.
- 7.2.1.4 Stockpile any existing topsoil during **landing** construction for later use in **rehabilitation**, this is not required if the operation uses suitable soil protection techniques (such as **cording** and **matting**).
- 7.2.2 Snig Track and Landing Rehabilitation
- 7.2.2.1 Following completion of the **timber harvesting operation** rehabilitate all **snig tracks** to prevent:
 - a) unacceptable movement of soil down or from the track surface; and
 - b) soil movement into streams.
- 7.2.2.2 Rehabilitate **landings** following completion of **timber harvesting operations**, and before the **coupe** is vacated, unless they are required for:
 - a) future Shelterwood 2 operations;
 - b) harvesting of adjacent coupes within 3 years; or
 - c) any other purpose authorised by the **Secretary** for which **native vegetation** is not compatible.

- 7.2.2.3 Identify any landings that do not require rehabilitation in the Forest Coupe Plan. 7.2.2.4 Lift and aerate corded and matted snig tracks to facilitate regeneration burning. 7.2.2.5 Remove cording and as much matting, bark and slash as possible from landings before rehabilitation works occur. 7.2.2.6 Rip / cultivate any snig track, landing or other area where machinery has compacted the soil. Ripping of **snig tracks** should be extended for at least 30 m from the **landing**. 7.2.2.7 Where removed and stockpiled, replace topsoil to a consistent depth across the landing. 7.2.3 **Boundary Trails** 7.2.3.1 Locate boundary trails within the gross coupe boundary and outside buffers, filters and exclusion areas except for sanctioned crossings identified in the Forest Coupe Plan. 7.2.3.2 Minimise debris deposited outside the **coupe** boundary when constructing boundary trails. 7.2.3.3 Maintain adequate drainage of boundary trails at all times until rehabilitation is complete. 7.2.3.4 Rehabilitate boundary trails as soon as practical after any regeneration burns and before commencement of any relevant closure periods. 7.2.4 Slash and bark management 7.2.4.1 Where **regeneration** burning is planned: a) place bark piles at least 10m inside the **coupe** boundary: b) place windrows at least 3m from excluded areas; and c) ensure slash is not permitted to accumulate within 3m of the base of any retained habitat tree, large tree or Shelterwood 1 tree. 7.2.4.2 Limit slash and bark piles to a maximum of 4 m² (ground area) and 10 m³ (total volume). 7.2.5 **Campsites and facilities** 7.2.5.1 Obtain approval from the Minister or delegate in accordance with Section 1.4 for any camp site or shower units associated with timber harvesting operations that are not located on a
- sanctioned coupe.
- 7.2.5.2 Situate camp sites and shower units in a location which requires no additional tree clearance.
- 7.2.5.3 In the Thomson, Tarago and Yarra Tributaries special water supply catchment areas serviced sanitary facilities must accompany any operation that is conducted on the land for any extended period.

7.3 Operational restrictions

7.3.1 Suspension of timber harvesting operations

- 7.3.1.1 Unless **cording** and **matting** has been established at a **coupe**, suspend **snigging**, **forwarding** and log loading when:
 - a) soil within the coupe is saturated;
 - b) water is flowing on or over any snig track; or
 - c) there is a reasonable risk that effective track drainage cannot be maintained.

7.3.2 Suspension of Haulage

- 7.3.2.1 **Haulage** must be suspended when:
 - a) snow is lying on any road used to access or exit the coupe;
 - b) water is flowing down any unsealed road or track; or
 - c) truck movement will deposit mud on a gravelled or sealed road.

7.3.3 Seasonal Closures

- 7.3.3.1 No harvesting or **stand tending** operations which involve the use of heavy machinery or which otherwise cause significant soil disturbance may take place in designated catchments in the Midlands **FMA** from 31 July to 31 October.
- 7.3.3.2 No **timber harvesting operations** may take place in the Woohlpooer and Cherrypool **State Forests** during the months of June, July, August and September.
- 7.3.3.3 Refer to Section 3.5 for season closures that apply to catchment areas within **SMZ**.

8. Fire Salvage Harvesting

8.1.1 Salvage General

- 8.1.1.1 Fire salvage prescriptions are in addition to all other prescriptions outlined in these **Management Standards and Procedures** and apply to all salvage operations within fire-affected **forest** until the beginning of the third winter following the wildfire, except where otherwise indicated.
- 8.1.1.2 **Timber harvesting operations** conducted within fire-affected **forest** after the third winter following the wildfire revert back to the non fire salvage prescriptions within these **Management Standards and Procedures**.
- 8.1.2 Salvage schedule priority
- 8.1.2.1 Undertake fire salvage harvesting operations in State forest areas designated as fire severity classes 1, 2 or 3 before salvage harvesting in fire severity classes 4 or 5. Table 24 Fire severity classification describes the Fire Severity Classification system.
- 8.1.3 Coupe size and aggregation
- 8.1.3.1 120 ha is the maximum **coupe** size for fire salvage operations in Alpine or Mountain **Ash** dominated **forest**. No size restrictions apply to aggregates of Alpine or Mountain **Ash** fire salvage **coupes**.
- 8.1.4 Habitat retention
- Plan the **coupe** so there is no more than 200 m between areas of retained habitat. Retained habitat includes **habitat tree exclusion areas**, filters and **buffers**, **green patch exclusion areas** and any **forest** adjacent to the **coupe**.
- 8.1.4.2 Exclude machinery from a minimum of 15 % of the gross coupe area to facilitate the recovery of understorey species. All exclusion areas, protection areas, SPZs and stream filters that are additional to standard stream protection prescriptions within the coupe may count towards the 15 % of gross coupe area from which machinery is excluded. 20 m is the minimum width for machinery exclusion areas set aside exclusively for understorey recovery. Where present, locate in areas with evidence of tree ferns. Tree felling is permitted in these areas. Cording and matting may remain on landings after salvage harvesting operations (replaces Clause 7.2.2.5).

8.1.5 Water quality

- 8.1.5.1 Except in **restricted access** and **special water supply catchment areas** locate **landings**, where possible:
 - a) at least 40 m for slopes of 15° or less, or 60 m for slopes of greater than 15°, from any **permanent stream**, or wetland; and
 - b) at least 20 m for slopes of 15° or less, or 30 m for slopes of greater than 15°, from any **temporary stream** and any **drainage line**.

- 8.1.5.2 In restricted access and **special water supply catchment areas** locate **landings**, where possible:
 - a) at least 60 m for slopes of 15° or less, or 80 m for slopes of greater than 15°, from any **permanent stream**, or **wetland**; or
 - b) at least 40 m from any temporary stream drainage line.
- 8.1.5.3 Locate boundary tracks at least 40 m from any **permanent stream** or **wetland**.
- 8.1.5.4 Apply the **buffer** and **filter** prescriptions for sites with high or very high water quality risk as specified in **Table 9 Minimum widths in metres for buffers (B) and filter (F) strips applicable to various waterway categories, in relation to water quality risk and slope.**
- 8.1.5.5 Install drainage structures for all **coupe** and **haulage** infrastructure at an appropriate frequency to mitigate increased risks to water quality due to potential sediment loads associated with water flow in the fire affected terrain.
- 8.1.5.6 In **restricted access** and **special water supply catchment areas** establish a drainage structure between 20 m to 40 m upslope of where a **road** or vehicle route crosses any **waterway**.

8.1.6 Weeds and disease

8.1.6.1 Clean soil from all harvesting machinery (excluding trucks and passenger vehicles) before floating to or from a salvage **coupe**. Replaces Clause 4.4.1.1 where pre-harvest disease and weed infestations cannot be assessed due to fire effects.

8.1.7 Field determined values

- 8.1.7.1 Locate the boundaries of **protection areas** or **management areas** based on modelled or mapped values according to the process outlined in **Figure 2 Field process for determining protection areas and management areas based on modelled or mapped values**.
- 8.1.7.2 To determine the boundaries of a Leadbeater's Possum habitat Zone 1A assume all **hollow bearing trees**, even if **dead**, to be 'alive'.

8.1.8 Ash Forest Green Patch Exclusion areas

- 8.1.8.1 Green patches are areas of **forest** within a fire affected area that are assessed through field checking to have fire severity class of 4 or 5 (See Section 8.1.2 and **Table 24 Fire severity classification**). Areas of fire severity class 4 or 5 must be at least 40 m wide to be a green patch.
- 8.1.8.2 Where less than 15 % of **Ash** dominated **forest** within a **compartment** is classified as fire severity class 4 or 5 all green patches greater than 0.5 ha are to be permanently protected during salvage operations to facilitate recruitment of future **hollow bearing trees**.
- 8.1.8.3 Where more than 15 % and less than 40 % of **Ash** dominated **forest** within a **compartment** is classified as fire severity class 4 or 5 protect all green patches greater than 5ha during salvage operations. Following salvage harvesting, further **timber harvesting operations** within these **compartments** are to be excluded until the **forest** regenerating from the fire reaches **reproductive maturity** typically by 15 years of age.
- 8.1.8.4 Green patches may be harvested where, within a **compartment**, more than 40 % **Ash** dominated **forest** is classified as fire severity class 4 or 5. Limit the total area of fire severity class 4 or 5 **Ash** harvested within a salvage **coupe** to 40 ha.

- 8.1.9 Ash habitat tree exclusion areas
- 8.1.9.1 Retain an average of at least 5 habitat trees per hectare of net coupe area in exclusion areas of greater than 0.1 ha. Replaces Table 12 Habitat tree prescriptions. Prescriptions protecting trees of pre-1900 origin continue to apply for Central Highlands FMAs.
- 8.1.9.2 Situate **habitat tree exclusion areas** to maximise retention of high priority **habitat trees**. **Habitat trees** have the following order of priority:
 - a) large live hollow bearing trees;
 - b) large live trees without hollows;
 - c) large dead trees;
 - d) small live trees; then
 - e) small dead trees.

Note: In the context of **salvage harvesting operations**, trees greater than 50 cm DBHOB are considered to be large. Replaces **habitat tree** selection criteria in Section 4.1, except where the Bendigo **FMA** standard applies or where all trees of a particular type (e.g. **hollow bearing / dead**) are protected.

- 8.1.9.3 Situate **habitat tree exclusion areas** to maximise retained **forest** connectivity within the **coupe**.
- 8.1.9.4 Arrange the shape and location of **habitat tree exclusion areas** to reduce ongoing operational and weather related damage to **habitat trees**.
- 8.1.10 Barred Galaxias buffers and filters
- Apply the minimum stream buffer and filter strip widths in Table 25 Salvage harvesting Barred Galaxia minimum buffer and filter strip widths upstream of Barred Galaxias populations (all soils), replacing Table 10 Minimum widths for buffer (B) and filter strips (F) applicable to waterway classes for specified aquatic and riparian habitat-dependant species.
- 8.1.10.2 Retain harvesting slash in **filter strips**, and aligned parallel to the stream, to slow the flow of water and reduce the potential for sediment to enter the stream or **wetland**.

9. Forest Regeneration and Management

9.1 Regeneration

- 9.1.1 Stocking surveys
- 9.1.1.1 For **even-aged stands**, conduct **stocking** surveys 15-30 months after sowing/seed fall or planting.
- 9.1.1.2 Successfully stocked **even-aged** regenerating **forests** must have:
 - a) at least 1 acceptable seedling in 65 % of 2.27 m radius (16 m²) plots placed on a 20 m by 80 m systematic grid or at least 1 acceptable seedling in at least 55 % of 2.27 m radius (16 m²) plots placed on a 20 m by 40 m systematic grid; and
 - b) no discrete area greater than 1 ha with less than 400 stems per hectare.
- 9.1.1.3 For **uneven-aged stands** conduct **stocking** surveys 15-36 months after completion of harvesting.
- 9.1.1.4 Successfully stocked **uneven-aged** regenerating **forest**s must have in at least 70 % of plots placed on a 20 m by 80 m systematic grid or 60 % of plots placed on a 20 m by 40 m systematic grid:
 - a) a total basal area of at least 30% of the basal area that the site is capable of supporting, or
 - b) at least 1 acceptable sapling (or coppice) in a 3.57 m radius (40 m²) plot or at least 1 acceptable sapling (or coppice) in a 3.57 m radius (40 m²) plot, or
 - c) at least 1 acceptable seedling in a 2.27 m radius (16 m²) plot or at least 1 acceptable seedling in a 2.27 m radius (16 m²) plot; and
 - d) no discrete area greater than 2 ha with less than 400 stems per hectare (caused by recent harvesting).
- 9.1.1.5 An acceptable seedling is defined as a species indigenous to the area in either a seedling, lignotuber, or coppice form. Seedlings, lignotubers and coppice must be over 40 cm in height for **Ash** species and over 25 cm in height for non **Ash** species. For coppice, the stem must be likely to remain attached to the stump and the base of the stem must be within 20 cm of ground level.
- 9.1.1.6 An acceptable sapling (can be in the coppice form) is defined as a species indigenous to the area that is an acceptable size (3–15 m), is not suppressed and has a vigorous, healthy **crown**.
- 9.1.1.7 For both **even-aged** and **uneven-aged silviculture**, acceptable species composition for **regeneration** includes at least 10 acceptable seedling of each eucalypt species present on the site before harvesting.
- 9.1.1.8 Where **stocking**, health or early growth is inadequate, remedial work must be conducted as soon as practicable and within 5 years of the previous **regeneration** attempt to obtain adequate **regeneration**. Further assessment must be undertaken following remedial treatment to ensure that it has been successfully regenerated.

- 9.1.2 Stocking Survey Exemptions
- 9.1.2.1 **Coupes** created purely for the purpose of **road construction** or **maintenance** do not require **stocking** surveys.
- 9.1.3 Seed
- 9.1.3.1 Prioritise seed collection from within areas available for **timber harvesting operations**.
- 9.1.3.2 Ensure large **hollow bearing trees** and **habitat trees** retained during previous **timber harvesting operations** are not felled for seed collection.
- 9.1.3.3 Collect seed from **stands** that show no evidence of hybridisation and have experienced a widespread and preferably heavy flowering in which trees with good crops are close together.
- 9.1.3.4 Collect seed only from **stands** of natural origin or artificially regenerated **stands** of satisfactory genetic status in which the full, original gene pool of the population is represented. **Stands** regenerated from **seed trees** are regarded as of natural origin for seed collection purposes.
- 9.1.3.5 Prioritise using seed collected from the **coupe**; i.e., regenerate with seed fall from **retained trees** and/or logging slash, or sow the **coupe** with seed collected from that **coupe**. Otherwise, prioritise using seed that meets the following criteria:
 - a) the collection site is within 25 km of the coupe to be sown;
 - b) the mid-elevation of the collection site is within 350 m above to 150 m below the midelevation of the **coupe** to be sown;
 - c) for slopes >10 ° the collection and sowing sites have similar aspect (separate between 'drier' aspects (W, NW, N, NE) and 'moister' aspects (SW, S, SE, E); and
 - d) the collection and sowing sites are of similar soil type and parent material.
- 9.1.3.6 Where seed quantities within areas available for **timber harvesting operations** are inadequate to meet **regeneration** requirements, application may be made to the **Minister** or **delegate** to collect seed from standing trees within areas excluded from **timber harvesting operations** in accordance with Section 1.4 and Clauses 9.1.3.7 to 9.1.3.10.
- 9.1.3.7 Applications to collect seed from areas excluded from **timber harvesting operations** will demonstrate that:
 - seed quantities from areas available for timber harvesting operations are inadequate to meet regeneration requirements;
 - b) trees will not be felled;
 - c) seed collection is on behalf of the Crown or the **Secretary**, or for **State forest regeneration** requirements;
 - d) there will be no machinery entry into rainforest and stream buffers;
 - e) protection area / SPZ values will be maintained; and
 - f) the operation will be assessed and monitored against any approval conditions.

- 9.1.3.8 Any **approved** seed collection operation in Leadbeater's Possum **SPZ** or associated **protection areas** will:
 - a) maintain canopy connectivity, particularly along roads;
 - b) protect understorey wattle from damage; and
 - c) exclude **hollow bearing** and pre-1900 aged trees from collection activities.
- 9.1.3.9 Applications to harvest seed from standing trees in areas excluded from harvesting will include:
 - a) the species to be collected;
 - b) an estimate of the extracted weight of seed to be collected; and
 - the FMA the seed will be collected from.
- 9.1.3.10 Submit applications at least 1 month prior to commencing seed collection.

9.2 Thinning

- 9.2.1.1 Unless otherwise stated, standards and procedures that apply to **timber harvesting operations** also apply to **thinning** operations.
- 9.2.1.2 In **Ash forests** retain at least 50 % of the live pre **thinning basal area** and a minimum total retained average **basal area** as outlined in **Table 26 Basal Area (BA) and stocking specifications for commercial ash thinning**.
- 9.2.1.3 In **mixed species forests** retain at least 50% of the live pre **thinning basal area** and a minimum total retained average **basal area** of:
 - a) 17 m2/ha for **stands** aged 18 24 years;
 - b) 21 m2/ha for **stands** aged 25 34 years; and
 - c) 25 m2/ha for **stands** aged 35+ years.
- 9.2.1.4 In **Box Ironbark forest**s retain at least 40 % of the live pre **thinning basal area** and a minimum total retained average **basal area** of 8 m2/ha (**regrowth** and **overwood**/retained habitat included).
- 9.2.1.5 In River Red Gum (*Eucalyptus camaldulensis*) **forest**s retain at least 40 % of the live pre **thinning basal area** and a minimum total retained average **basal area** of 15 m2/ha in riverine and 10 m2/ha in dryland **stands** (regrowth and **overwood**/retained **habitat trees**).

Appendix 1: Tables and Figures

Table 1 Soil susceptibility to breakdown

					Horizon	
Soil factor		Scoring		Α	В	C
Soil texture Score	SC, C, SiC 1	SiCL, CL, SCL 4	SL, L, SiL, LS, S, Si 7			
Soil aggregate stability Score	Unaffected 1	Slaking, slight dispersibility 7	Very dispersible 15			
Soil structure Score	Highly pedal 1	Peds evident 3	Apedal 5			
Soil colour	Black, Reddish Black, Dark Brown 1	Light Brown, Light Red 2	Pale Yellow, Grey, White 3			
Soil organic matter	High 1	Moderate 4	Low 7			
Stoniness Score	>50% 1	10-50% 2	<10% 3			
Overall score	<17	17-27	>27			
Rating	Low	Moderate	High			

Table 2 Soil permeability

					Horizon	
Soil factor		Scoring		Α	В	С
Soil texture Score	SL, L, LS, S 1	SiL, Si, SiCL, CL, SCL 4	SC, C, SiC			
Organic matter	High	Moderate	Low			
Score	1	2	3			
Impeding layer	>50	50-25	<25			
depth (cm) Score	1	3	5			
Soil Mottling	Not present	Slight	Well Developed, (Pale)			
Score	1	2	(1 ale)			
Stoniness	>50%	10-50%	10%			
Score	1	2	3			
Stand ht (m)	>40	28-40	<28			
Score	1	2	3			
Overall score	<11	11-18	>18			
Rating	Very high	High	Low			

Table 3 Soil erosion site factor

Site factor		Scoring	
Erosivity index	<590	591-1469	>1470
Rating	1	3	5
Slope (degrees)	0-8	9-21	>21
Rating	2	4	7
Slope length / uniformity	Short uniform	Long broken	Long uniform
Rating	1	2	3
Revegetation capacity	Good	Moderate	Poor
Rating	1	2	3
Overall score	<10	10-14	>14
Rating	Low	Moderate	High

Notes:

- Assess soil when moderately moist.
- Stoniness refers to stones greater than 20 mm in size.
- Stand height refers to a mature forest stand height.




Figure 1 Rainfall erosivity

Rainfall erosivity contours in non-SI units for Victoria (Garvin et al. 1979). SI units (MJ. Mm/ha.h.year) given in parentheses.

Table 4 Soil erodibility classification

	Table 1 rating				•		
Table 2 rating	Low	Moderate	High	Horizon	Α	В	С
Very High	VERY LOW	LOW	MEDIUM	Table 1			
High	LOW	MEDIUM	HIGH	Table2			
Low	MEDIUM	HIGH	VERY HIGH	Classification			

Table 5 Soil erosion hazard classification

Table 4 classification									
Table 3 rating	Very low	Low	Medium	High	Very high	Horizon	A	В	С
Low	LOW	LOW	LOW	MEDIUM	HIGH	Table 4			
Moderate	LOW	LOW	MEDIUM	HIGH	VERY HIGH	Table 3			
High	LOW	MEDIUM	HIGH	VERY HIGH	VERY HIGH	Classification			

Table 6 Soil permeability classification

	Soils with HIGH permeability	Soils with LOW permeability	Horizon	Α	В	С
Table 2 score	0 - 18	19+	Table 2			
			Classific	ation		

Table 7 Overall soil category

Soil Permeability	Soil Erosion Classification (Table 5)						
Classification (Table 6)	Low	Medium	High	Very High			
Low	М	М	Н	VH			
High	L	L	M	М			

Table 8 Water quality risk

Overall Soil Category	Water Quality Risk	
L	L Low water quality risk	
М	Moderate water quality risk	
Н	High water quality risk	
VH	Very High water quality risk	

Table 9 Minimum widths in metres for buffers (B) and filter (F) strips applicable to various waterway categories, in relation to water quality risk and slope

	Sites with low or moderate water quality risk	Sites with high or very high water quality risk		
Waterway Class	Slope 0 - 30 °	Slope 0 - 20 °	Slope 21 - 30 °	
Pools, permanent streams and wetlands	20 m B	30 m B	40 m B	
2. Temporary streams	10 m F	10 m B + 10 m F	20 m B	
3. Drainage lines	10 m F	10 m F	15 m F	

Table 10 Minimum widths for buffer (B) and filter strips (F) applicable to waterway classes for specified aquatic and riparian habitat-dependant species

	Sites with low or moderate water quality risk	Sites with high or very high water quality risk		
Waterway Class	Slope 0-30°	Slope 0-20°	Slope 21° - 30°	
Pools, permanent streams and wetlands	30 m B	40 m B	50 m B	
2. Temporary streams	20 m B + 10 m F	20 m B + 20 m F	30 m B + 20 m F	
3. Drainage lines	10 m F	10 m F	15 m F	

Table 11 Water supply protection areas

Catchment	Status	Slope Limit	Seasonal Closure	Stream Buffers	Filter Strips	Additional Management Actions or Comments
Benalla - Mans	field FMA					
Lake Eildon	SWSCA	30°	1 June - 31 Oct	20 m	10 m	
Upper Goulburn	SWSCA	30°	1 June - 31 Oct	20 m	10 m	
Lake Nillahcootie	SWSCA	30°	1 June - 31 Oct	20 m	10 m	
Ryans Creek	SWSCA; DC					Max 40 ha max coupe size
	Category 1:	-	-	-	-	Category 1: No harvesting
	Category 2:	< 12 °	1 May - 31 Oct	20 m	10 m	Category 2: No clearfall
	Category 3:	30°	1 May - 31 Oct	20 m	10 m	Category 3: NA
Central FMA						
Sunday Creek (Broadford- Kilmore)	SWSCA	30°		20 m	10 m	
Kilmore	SWSCA	30°		20,40, 60 m	10 m	Buffers in accordance with former <i>Notice</i> of <i>Determination</i> of <i>Land Use</i> , <i>Plan No.</i> 1633.
Upper Goulburn	SWSCA	30°		20 m	10 m	
Dandenong FM	IA ²					
Bunyip River	SWSCA	25°	1 May - 30 Nov	20 m	10 m	60 m buffer upstream of weir.

Britannia Creek	SWSCA	30°	1 July – 30 Sept	20 m	10 m	
Tomahawk Creek (Gembrook)	SWSCA	30°	1 July – 30 Sept	20 m	10 m	
McCrae's Creek	SWSCA	30°	1 July – 30 Sept	20 m	10 m	60 m buffer upstream of weir.
Micks Creek	SWSCA					Harvesting excluded.
Healesville	SWSCA					Harvesting excluded.
Armstrong Creek East	Restricted					Harvesting excluded.
Learmonth Creek		25°	1 May – 30 Nov	40 m ¹	10 m	Max 7 ha Ash and 3ha mixed species .
McMahons Creek	Restricted	25°	1 May – 30 Nov	40 m ¹	10 m	A maximum of 30 % of each of these catchments is to be harvested over a 10
Starvation Creek	Restricted	25°	1 May – 30 Nov	40 m ¹	10 m	year period. There is to be harvesting in only 1 of the 4 restricted access
Cement Creek	Restricted	25°	1 May – 30 Nov	40 m ¹	10 m	catchments in any 1 year.
Armstrong Creek West	Restricted	25°	1 May – 30 Nov	40 m ¹	10 m	-
East Gippsland	FMA					
Orbost (Rocky River)	SWSCA			100m 40m 30m 100m		40 ha maximum annual area harvested. Buffer applies to main river Buffer applies to major tributaries Buffer applies to other watercourses Buffer applies to off-take weir.
Betka River	SWSCA			150m		120 ha (1% of catchment) max annual area harvested On main river
0'	. 3			20m		On major tributaries and streams
Gippsland FMA Thomson	SWSCA	30°	1 May – 30 Nov	40 m ¹	10 m	Subject to additional regulations
River	SWSCA	30	Tiviay – 30 Nov	40 111	10 111	under the Forests Act 1958 which includes limitations on access, camping, hunting with hounds and horse riding. Average 150 ha Ash and 15ha mixed species on rolling average, or total of 165 ha on rolling average.
Tyers River	SWSCA	30°	1 June – 31 Oct	20 m	10 m	Applies above 650 m ASL. Harvesting restrictions on land above 1220 m. Buffer within 200 m of Moondarra Reservoir foreshore. Buffer of 40 m on Jacobs Creek up to the Walhalla Road.
Tanjil River	SWSCA	30°	1 June – 31 Oct	20 m	10 m	Closure applies north of Icy Creek – Mt Baw Baw Road.
Tarago River	SWSCA	30°	1 May – 31 Oct	20 m	10 m	Average 55 ha Ash and 23 ha mixed species on rolling average.
Bunyip River	SWSCA	25°	1 June – 30 Nov	20 m	10 m	
Deep Creek and Loch River (Noojee)	SWSCA	30°	1 July – 30 Sept	20 m	10 m	
Drouin	SWSCA	30°	1 July – 30 Sept	20 m	10 m	
Glenmaggie	SWSCA		1 June – 31 Oct	40 m		Buffer applies to Macalister below Caledonia
				40 m		Buffer applies to Wellington below Carey
				40 m		Buffer applies to Barkley below Mt Skene Creek
				40 m		Buffer applies to Glenmaggie Creek below east and west branches
Mirboo North	SWSCA			Buffer of	of 100 m ai	round the off-take weir

 ${\bf Buffer}$ from the banks of streams, watercourses and spring areas. 40 m ${\bf buffer}$ on the Little Morwell River and 20 m for other water courses.

				20 m for other water courses.
Tarra River	SWSCA		1 June – 31 Oct	
North East FMA				
Lake Hume (Victorian section)	SWSCA		1 July – 30 Sept	
Mount Tabor Creek	DC		1 July – 30 Sept	40 ha maximum annual area harvested.
Black Dog Creek	DC		1 July – 30 Sept	30 ha maximum coupe size.
Bakers Gully (Bright)	SWSCA; DC		1 July – 30 Sept	20 ha maximum coupe size.
West Kiewa River	DC		1 July – 30 Sept	40 ha maximum coupe size.
Musk Gully Creek	DC		1 July – 30 Sept	35 ha maximum coupe size.
Diddah Diddah Creek (Springhurst)	SWSCA; DC		1 July – 30 Sept	20 ha maximum coupe size.
Midlands FMA				
Blackwood	DC		1 June – 31 Oct	35 ha maximum coupe size.
Bullarto	DC		1 June – 31 Oct	20 ha maximum coupe size.
Colbrook	DC		1 June – 31 Oct	5 ha maximum coupe size.
Collier Gap	DC		1 June – 31 Oct	2 ha maximum coupe size.
Djerriwarrh	SWSC & SAP; DC		1 June – 30 Sep	15 ha maximum coupe size.
Korweingu- boora	DC		1 June – 31 Oct	15 ha maximum coupe size.
Hickmans Creek (Elmhurst)	DC		1 June – 31 Oct	40 ha maximum coupe size.
Lal Lal Lake Environs	SWSC & SAP; DC		1 June – 31 Oct	5 ha maximum coupe size.
Long Gully	DC		1 June – 31 Oct	15 ha maximum coupe size.
Lake Merrimu	SWSC & SAP; DC		1 June – 30 Sep	40 ha maximum coupe size. Buffers apply to all waterways
Moorabool	DC		1 June – 30 Sep	10 ha maximum coupe size.
Mount Cole	DC		1 June – 31 Oct	20 ha maximum coupe size.
Musical Gully	SWSC; DC		1 June – 30 Sep	2 ha maximum coupe size.
Pykes Creek	SWSC; DC		1 June – 30 Sep	40 ha maximum coupe size.
Rosslynne	SWSC & SAP; DC		1 June – 30 Sep	40 ha maximum coupe size.
Shepherds Creek	DC		1 June – 31 Oct	40 ha maximum coupe size.
Sugarloaf	DC DC		1 June – 30 Sep	20 ha maximum coupe size.
Troy	SWSC; DC		1 June – 30 Sep	2 ha maximum coupe size.
White Swan	DC		1 June – 31 Oct	20 ha maximum coupe size.
Wombat	DC		1 June – 31 Oct	20 ha maximum coupe size.
Otway FMA ³		250	1 May 20 Na	May harvoot 0.5.0/ new year and 4.5.
Lorne – St Georges River	CMCCA	25°	1 May – 30 Nov	Max harvest 2.5 % per year and 15 % per decade of total forested area of each catchment.
Barham River	SWSCA	25°	1 May – 30 Nov	
Upper Barwon	SWSCA	25°	1 June – 31 Oct	Max harvest 5 % per year and 25 % per decade of total forested area of
Gellibrand River	SWSCA	25°	1 June – 31 Oct	each catchment.
Painkalac Creek (Aireys Inlet)	SWSCA	25°	1 June – 30 Sept	
Pennyroyal Creek	SWSCA	25°	1 June – 30 Sept	

Matthews Creek	SWSCA	25°	1 June – 30 Sept	
Gosling Creek	SWSCA	25°	1 June – 30 Sept	
Tambo FMA				
Lake Hume (Victorian section)	SWSCA		30 June – 1 Oct	In accordance with former Plan No. S-1275B.
Mitchell River	SWSCA		30 June – 1 Oct	Seasonal closure applies above 900 m.
Nicholson River	SWSCA	30°		200 m buffer around the Nicholson Reservoir and Water supply off-take on the Nicholson River. 40 m buffer around Nicholson and Barmouth Rivers. 20 m buffer from banks of permanent stream s & drainage lines

Notes:

- Restricted restricted access catchments managed by agreement between Melbourne Water and the **Department**.
- ¹ Major streams only.
- ² In addition to the above prescriptions for the **Gippsland FMAs** and the Dandenong **FMA**, refer to clauses 3.5.1.5 and 3.5.1.6 of these **Management Standards and Procedures**.
- ³ Within the Otway **FMA**, 20° and 15° slope limits will be applied in **special water supply catchment areas** in areas of Land Degradation Hazard Class 4 and 5 respectively.

Table 12 Habitat tree prescriptions

Locality	Forest Type	Habitat Tree Retention Rates	Comment
Central Highlands FMAs	Ash /HEMS	All live ash eucalypts originating before 1900. At least 40 trees per 10 ha for the length of the rotation in ash forests originating since 1900	Retain at least 1 potential hollow bearing tree where gaps between retained trees are greater than 150 meters, Retained trees should be a mixture of hollow bearing trees where present and other trees most likely to develop hollows in the short term.
	Mixed Species	40+ trees per 10 ha	
East Gippsland FMA	All	4 – 5 trees per ha	Count seed trees towards habitat tree numbers.
Gippsland FMAs	All	4 – 5 tree per ha	Count seed trees towards habitat tree numbers
Horsham FMA	All	10 trees per 10 ha 5 potential habitat trees per 10 ha 5 dead trees with hollows per 10 ha	Any additional trees above the 10 trees per 10 ha should be retained in place of potential habitat and dead trees.
Midlands FMA	All except Box Ironbark	3 trees per ha	Refer to Bendigo for Box ironbark prescriptions.
Mid Murray FMA	All	20 trees per 10 ha (≥50 cm but ≤100 cm DBHOB) Retain all trees (>100 cm DBHOB)	
North East FMAs	Ash/HEMS	4 – 5 trees per ha	
	Mixed species	4 – 5 trees per ha	Count seed trees towards habitat tree numbers.
Otway FMA	All	5 trees per ha (net coupe area) / 10 trees per ha (gross coupe area)	
Portland FMA	All	Retain habitat trees to a basal area of 2 – 4m²/ha	Group selection.

Table 12 Notes:

- In all cases except the Otway FMA, habitat tree retention rates apply relative to the net harvested area retained within the net coupe area.
- The **net coupe area** is the **gross coupe area**, less areas required to be excluded from timber harvesting in accordance with the **Code**.
- Areas excluded from timber harvesting, including **buffers**, should be identified prior to the commencement of **timber harvesting operations**, roading and regeneration.
- Areas excluded from timber harvesting or retained areas may not be readily identifiable prior to timber harvesting operations but must be documented after timber harvesting operations have been completed (e.g. small rocky areas, small areas of steep slope or inaccessible areas), these areas may be counted as retained habitat within the **net coupe area**.
- The minimum number of trees required to be retained as habitat relative to the area harvested (net harvested area) is a product of the specified rate and the **net harvested area**. E.g., 20 ha **net harvested area** × 4 - 5 trees per ha = 80 - 100 trees to be retained has habitat.
- This total number of trees retained as habitat can be dispersed across the **coupe** or retained in clumps, prioritising the types and arrangement of trees as relevant to section 4.1 of the MSP.

Table 13 Rare or threatened fauna and invertebrate prescriptions

Species Name	Value	Applicable FMAs	Management Actions
Alpine She-oak Skink Cyclodomorphus praealtus	Individual (site occupied by at least 1 observed individual)	North East FMAs	Develop management actions to protect Alpine She-oak Skink habitat at sites occupied by at least 1 observed individual. Document and provide to the Secretary a proposed plan for the value management area .
	Individual (site occupied by at least 1 observed individual)	Gippsland FMAs	Apply a protection area of approximately 50 ha over each site occupied by at least 1 observed individual.
Alpine Tree Frog Litoria verreauxii alpina	Population	North East FMAs	Apply a management area over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations . Document and provide to the Secretary a proposed plan for the value management area.
Alpine Water Skink Eulamprus kosiuskoi	Population	Gippsland FMAs North East FMAs	Apply a management area over populations. (management area conditions - Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations).
Apostlebird Struthidea cinerea	Population	North East FMAs	Apply a management area over sites of resident populations. (management area conditions - Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations).
Australasian Darter Anhinga novaehollandiae	Roosting and breeding site	Mid-Murray FMA	Exclude activities likely to disturb breeding activity within 250 m region around current roosting and breeding sites of colonially-nesting waterbirds during the breeding season
Australian Grayling Prototroctes maraena	Population	Portland-Horsham FMA East Gippsland FMA	Apply a protection area extending 100 m from each bank for 1 km upstream and 1 km downstream of populations. Avoid constructing new roads in the protection area . Note: The Secretary intends to review this strategy when 50 significant fish sites have been located.
Australian Hobby Falco longipennis	Nest tree	Otways FMA	Apply a protection area of 100 m radius around nest trees.
Australian White Ibis	Roosting and breeding site	Mid-Murray FMA	Exclude activities likely to disturb breeding activity within 250 m region around current roosting and

Species Name	Value	Applicable FMAs	Management Actions
Threskiornis molucca			breeding sites of colonially-nesting waterbirds during the breeding season
	Individual (site occupied by at least 1 observed individual)	North East FMAs	Apply a protection area of approximately 50 ha over each site, where practicable incorporating the detection site.
Bandy Bandy	oboorvod individualy		Note: The Secretary intends to review the strategy when 20 SPZ are established.
Vermicella annulata			Apply a management area of approximately 50 ha over sites, where practicable incorporating the detection site.
	Individual (site occupied by at least 1 observed individual)	Mid-Murray FMA	Conduct a site inspection and detailed planning in consultation with DELWP to ensure the species is adequately protected during timber harvesting operations .
	Roosting and nesting tree	Statewide	Apply a protection area of 3 ha and apply a management area of 250-300 m radius (or equivalent linear area) around each nesting and roosting site utilised recently and frequently and located outside a Barking Owl Management area .
			(A management area is not required where existing zoning or public land tenure provides equivalent or greater protection of the roosting or nesting site).
Barking Owl Ninox connivens			Document and provide to the Secretary a proposed plan for the value management area prior to harvesting commencement. The plan will protect hollow bearing trees for owls and prey species along with younger trees that may provide hollows in future. These areas will be managed to maintain habitat capable of supporting adequate populations of birds and arboreal prey mammals to support breeding owls. Exclude timber harvesting operations in the management area or SMZ s around nest trees during the breeding season (1 July - 31 December).
Barred Galaxias Galaxias fuscus	SMZ for Barred Galaxias or waterways containing	Central Highlands FMAS Gippsland FMAs	Apply minimum stream buffer and filter strip widths (specified in Table 10) 1km upstream of new Barred Galaxias populations or in management areas / SMZ for Barred Galaxias.
	new population	North East FMAs	Minimise stream crossings over waterways in catchments containing Barred Galaxias.

Species Name	Value	Applicable FMAs	Management Actions
Black Falcon Falco subniger	Nesting tree	Portland-Horsham FMA	Apply a management area of 250 m radius over each nesting site. Within 250 m of nesting site, avoid timber harvesting, road construction and burning during breeding season. Apply a protection area with a 100 m radius around nest trees
Blue Mountains Tree Frog Litoria citropa	Population of more than 50 individuals	East Gippsland FMA	Apply a protection area extending 100 m from each bank for 500 m upstream and 500 m downstream for sites of more than 50 individuals. Note: The Secretary intends to review this strategy when 20 sites are established.
Brush-tailed Phascogale Phascogale tapoatafa	Individual (site occupied by at least 1 observed individual) in areas where species is not understood to occur	Statewide	Within FMAs where the species is not understood to occur, notify the Secretary and determine whether the value occurs in a Brush-tailed Phascogale Management area (BPMA) or whether a new BPMA may be required. If a new BPMA is required, apply a management area that protects and maintains the Brush- tailed Phascogale population and their habitat, retaining large trees and naturally fallen timber. Note: The Secretary intends to review and change sites only if new sites are of a higher quality or are better able to be protected than existing sites.
Bush Stone- curlew Burhinus grallarius	Nesting site	Portland-Horsham FMA	Apply a management area of 250 m radius over each nesting site. Within the management area, avoid any burning during breeding season. At all other times retain any logs and other fallen timber in the area.
	Population	Mid Murray FMA	Protect populations from disturbance where practicable during the hibernating (1 April to 30 September) and breeding (1 November to 28 February) periods respectively.
Carpet Python Morelia spilota metcalfei	Population	North East FMAs	Apply a management area of 250 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Common Bent- wing Bat Miniopterus	Colony	Central Highlands FMAs	Apply a 100 m buffer around colonies.

Species Name	Value	Applicable FMAs	Management Actions
schreibersii oceanensis		North East FMAs	
			Apply a management area of 100 m radius over roosting and over-wintering sites.
	Roosting and over- wintering site	Midlands FMA	Within 100 m of roosting and over-wintering sites, exclude timber harvesting operations, road construction and other activities likely to disturb overwintering bats between 1 May to 31 October.
	Breeding and roosting cave, tree, mine or over-wintering site	Gippsland FMAs East Gippsland FMA	Apply a protection area of 100 m radius over breeding and roosting caves, trees and mines and over-wintering sites.
Common Brush- tailed Possum Trichosurus vulpecula	Relative abundance (More than 10 per Spotlight Kilometre)	East Gippsland FMA	Apply a protection area of approximately 100 ha of suitable habitat where records report a relative abundance of more than 10 individuals per spotlight kilometre (equivalent to more than 2 individuals per hectare or more than 15 individuals per hour of spotlighting), or where substantial populations are located in isolated or unusual habitat.
vuipecuia			Note: Assumed rate of spotlighting per kilometre is 100 mins per 1km and visible range either side of transect for this species is 25m, equating to assumed minimum survey area of 5 hectares.
Common Dunnart	Population	Central Highlands FMAs Midlands FMA Portland-Horsham FMA	Apply a protection area of approximately 20 ha of suitable habitat over sites.
Sminthopsis murina murina			Note: The Secretary intends to review this strategy when 50 sites have been established in State forest .
Cox's Gudgeon			Apply a protection area extending 100 m from each bank for 1 km upstream and 1 km downstream of populations.
Gobiomorphus coxii	Population	East Gippsland FMA	Avoid constructing new roads in the protection area.
			Note: The Secretary intends to review this strategy when 50 significant fish sites have been located.
Diamond Python	5 1.0	O:	Apply a protection area of 100 ha of suitable habitat for each locality of this species.
Morelia spilota spilota	Population	East Gippsland FMA	Note: The Secretary intends to review this strategy when 50 sites have been located.
Dwarf Galaxias Galaxiella pusilla	Population	Portland-Horsham FMA	Minimise stream crossings over permanent and temporary streams and drainage lines in catchments containing populations of Dwarf Galaxias.
Eastern Broad- nosed Bat Scotorepens orion	Population Or	East Gippsland FMA	Apply a protection area of approximately 100 ha of older forest over sites where more than 10 individuals have been captured in one trapping session.

Species Name	Value	Applicable FMAs	Management Actions
	Relative abundance per Spotlight Kilometre		Note: The Secretary intends to review this strategy when 20 sites are established.
Eastern Great Egret Ardea modesta	Current roosting and breeding site	Portland-Horsham FMA	Exclude activities likely to disturb breeding activity within 250 m region around current roosting and breeding sites of colonially-nesting waterbirds during the breeding season.
Eastern Horseshoe Bat Rhinolophus	Colony	Central Highlands FMAs	Apply a 100 m buffer around colonies.
megaphyllus megaphyllus	Breeding or roosting cave, mine, tree, or over-wintering site	Gippsland FMAs East Gippsland FMA	Apply a protection area of 100 m radius over breeding and roosting caves, trees and mines and over-wintering sites.
Eastern Pygmy- possum Cercartetus nanus	Relative abundance (More than 5 per pitfall trap)	East Gippsland FMA	Where records report more than 5 individuals per standard pitfall line over 5 days, or where substantial populations are in isolated or unusual habitat, apply a protection area of approximately 100 ha of suitable habitat.
Eltham Copper Butterfly Paralucia pyrodiscus lucida	Population Host plant <i>Bursaria</i> <i>spinosa</i>	Portland-Horsham FMA	Apply a management area over populations. Protect the host plant Bursaria spinosa (Sweet Bursaria) within 250 m of any records of this species.
Four-toed Skink Hemiergis peronii	Population	Midlands FMA	Apply a management area over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Freshwater Herring Potamalosa richmondia	Population	East Gippsland FMA	Apply a protection area extending 100 m from each bank for 1 km upstream and 1 km downstream of populations. Avoid constructing new roads in the protection area . Note: The Secretary intends to review this strategy when 50 significant fish sites have been located.
Giant Burrowing Frog Heleioporus australiacus	Population	Gippsland FMAs East Gippsland FMA	Where records of Giant Burrowing Frog are located on first-order streams or sites away from streams, apply a protection area of 50 ha (preferably the entire sub catchment unit). Where records of Giant Burrowing Frog are located on

Species Name	Value	Applicable FMAs	Management Actions
			second or higher order streams, apply a protection area of 100 m width each side of the stream for 1km upstream and downstream of the detection site.
			Note: For the purposes of this prescription, a first order stream is the headwaters of a catchment and is the smallest stream mapped on the 1:100 000 Natmap series. Second order streams are the next level of stream further down the catchment. For first order streams the size of each catchment will vary, however target size is approximately 50 ha.
			For off stream records where evidence of this value is found in the field, apply a protection area of 50ha over the record or equivalent area of suitable habitat nearby.
			Avoid new roading in the protection area .
			Note: The Secretary intends to review this strategy when 50 sites are located in Victoria.
Giant Gippsland			Apply a management area over populations.
Earthworm Megascolides australis	Population	Gippsland FMAs	Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Glenelg Freshwater Mussel	Population	Portland-Horsham FMA	Apply a protection area extending 100 m from each bank for 1 km upstream and 1 km downstream of populations.
Hyridella glenelgensis			Disturbance that could impact on water quality must be avoided within the protection area .
			Apply a management area of 250 m radius over each nesting site.
Glossy Black- Cockatoo		Gippsland FMAs	Within 250 m of nest tree, exclude timber harvesting operations , road construction and burning during the breeding season (December to May). Search the surrounding forest for other active nest sites (the species is known to nest in clusters). Apply a protection area with a 100 m radius around nest trees.
Calyptorhynchus lathami lathami	Nesting site	East Gippsland FMA	Where Black She-oak stands are identified during timber harvesting operations (including post harvesting burning) new roading activities will be conducted in a manner that avoids damage to the stand .
			The definition of a Black She-oak stand is a group or groups of trees with a basal area equal to or greater than 10 m2 in an area of 0.25 hectares.

Species Name	Value	Applicable FMAs	Management Actions
Golden Sun Moth Synemon plana	Population	Portland-Horsham FMA	Apply a management area over populations. Detailed planning is required to ensure population is protected from timber harvesting.
Great Cormorant Phalacrocorax carbo	Population	Mid-Murray FMA	Exclude activities likely to disturb breeding activity within 250 m region around current roosting and breeding sites of colonially-nesting waterbirds during the breeding season.
Greater Glider Petauroides volans	Relative abundance (More than 10 per Spotlight Kilometre)	East Gippsland FMA	Apply a protection area of approximately 100 ha of suitable habitat where records report a relative abundance of more than 10 individuals per spotlight kilometre (equivalent to more than 2 individuals per hectare or more than 15 individuals per hour of spotlighting), or where substantial populations are located in isolated or unusual habitat. Note: Assumed rate of spotlighting per kilometre is 100mins per 1km and visible range either side of transect for this species is 25m, equating to assumed minimum survey area of 5 hectares.
	Nest tree used within the last 5 years	Central Highlands FMAs North East FMAs	Apply a 100 m buffer around nest trees used within the last 5 years. Exclude timber harvesting operations within 250 m of nest trees during breeding season.
Grey Goshawk	Nest tree	Otways FMA	Buffer nest trees by 100 m and avoid burning at all times.
Accipiter novaehollandiae	Nesting sites	Midlands FMA Gippsland FMAs Portland-Horsham FMA Bendigo FMA East Gippsland FMA	Apply a management area of 250 m radius over nesting sites. Within 250m of nesting site, avoid timber harvesting, road construction and burning activities during breeding season. Breeding season is 1 July to 30 November. Apply a protection area of 100m radius around nest trees. Note: The Secretary intends to review this strategy when 10 sites have been established in State forest.
Grey-crowned Babbler Pomatostomus temporalis	Active colony	Mid-Murray FMA	Apply a buffer of 200 m radius around active colonies utilising River Red Gum (<i>Eucalyptus camaldulensis</i>) forest on the western edge of Guttram State forest and the southern edge of Benwell State Forest . Note: The Secretary intends to review this prescription when stand type and structural habitat requirements are understood. Establish a buffer of 100 m radius around active colonies elsewhere.
Grey-headed Flying-fox Pteropus poliocephalus	Roost site that has a pattern of regular seasonal use Active colony	Gippsland FMAs	Apply a protection area of 100 m radius for a roost site that has a pattern of regular seasonal use. Schedule timber harvesting operations and burning
	, 10.110 001011y		

Species Name	Value	Applicable FMAs	Management Actions
			to minimise disturbance to active colonies if found in State forest.
	Active colony	Portland-Horsham FMA	Schedule timber harvesting operations to minimise disturbance to active colonies if found in State forest .
Hooded			Apply a management area of approximately 50 ha over populations, where practicable incorporating the detection site.
Scaly-foot Pygopus schraderi	Population	Mid-Murray FMA	Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Plumed Egret Ardea intermedia	Current roosting and breeding site	Portland-Horsham FMA	Exclude activities likely to disturb breeding activity within a 250 m radius of current roosting and breeding sites of colonially-nesting waterbirds during the breeding season.
plumifera	Current roosting and breeding site	Mid-Murray FMA	Exclude activities likely to disturb breeding activity within a 250 m radius of current roosting and breeding sites of colonially-nesting waterbirds during the breeding season
King Quail Synoicus chinensis	Wet heath	Portland-Horsham FMA	Protect wet heaths within 20ha radius of records of this species. Harvesting should be minimised close to wet heaths and drainage patterns must not be altered.
Koala Phascolarctos cinereus	Resident population or substantial population located in isolated or unusual habitat	East Gippsland FMA	Apply a protection area of approximately 100 ha of suitable habitat over resident populations or substantial populations located in isolated or unusual habitat.
Lace Monitor Varanus varius	Population	Portland-Horsham FMA	Apply a management area over populations. Detailed planning is required to ensure population is protected from timber harvesting.
Large Brown Tree Frog Litoria littlejohni	Detection (adult, sub- adult, tadpole or egg cluster)	East Gippsland FMA	Apply a protection area of 28 ha that includes the detection site.
Leadbeater's Possum habitat/colony	Leadbeater's Possum colony OR	Central Highlands FMAs	Colony Apply a protection area of 200 m radius centred on each Leadbeater's Possum colony.

Species Name	Value	Applicable FMAs	Management Actions
Gymnobelideus leadbeateri	Zone 1A habitat		OR
leaubeateri	OR		Zone 1A Habitat
	Zone 1B habitat		Apply a protection area over areas of Zone 1A habitat where there are more than 10 hollow bearing trees per 3 ha in patches greater than 3 ha. (Ensure Zone 1A habitat is not salvage logged).
			Zone 1B Habitat Apply a protection area over Zone 1B habitat where there are more than 12 hollow bearing trees per 3 ha in patches greater than 10 ha and wattle density exceeds 5 m2/ha. This prescription applies until either of the two Zone 1B attributes:
			1. the presence of dead mature or senescent living trees; or
			2. wattle understorey no longer exists.
Little Black Cormorant Phalacrocorax sulcirostris	Current roosting and breeding site	Mid-Murray FMA	Exclude activities likely to disturb breeding activity within 250 m radius of current roosting and breeding sites of colonially-nesting waterbirds during the breeding season.
Little Egret Egretta garzetta	Current roosting and breeding site	Portland-Horsham FMA	Exclude activities likely to disturb breeding activity within 250 m radius of current roosting and breeding sites of colonially-nesting waterbirds during the
Lgrona garzona		Mid-Murray FMA	breeding season.
			Apply a management area of 250 m radius over each nesting site.
Little Falcon Falco longipennis	Nesting site	East Gippsland FMA	Within 250 m of nesting site, avoid timber harvesting, road construction and burning during breeding season. Apply a protection area of 100 m radius around nest trees.
Little Pied Cormorant Microcarbo melanoleucos	Current roosting and breeding site	Mid-Murray FMA	Exclude activities likely to disturb breeding activity within 250 m radius of current roosting and breeding sites of colonially-nesting waterbirds during the breeding season.

Species Name	Value	Applicable FMAs	Management Actions
Long-footed Potoroo Potorous longipes	Detection site	North East FMAs East Gippsland FMA	Apply a management area of approximately 150 ha for each Long-footed Potoroo detection site that is outside the Core Protected Area (i.e. within SPZ and conservation reserves). Notes: As far as possible, management area boundaries will follow recognisable landscape features such as ridges, spurs and watercourses. Within each management area, at least one third (50 ha) will be protected from timber harvesting and new roading. This will be known as Long-footed Potoroo Retained Habitat. The LFP Retained Habitat will include the best LFP habitat in the SMZ, which will generally be in gullies and on lower, sheltered slopes. The LFP Retained Habitat may include areas otherwise unavailable for timber harvesting due to restrictions under the Code. The management area will also have a general restriction of one third of the total area that can be harvested in any three year period. If more than one coupe is to be harvested in a management area in the same year, the coupes must be separated by at least the equivalent of another coupe width. The management area, with the LFP Retained Habitat clearly delineated, will be shown as part of the Forest Management Zoning Scheme. If the 150 ha area includes any part of an existing conservation reserve or SPZ, these areas will retain their existing reservation or zoning status but will be considered for inclusion as part of the area of retained habitat. In such cases, the final area applyed as a management area or SMZ may be correspondingly smaller.
Long-nosed Potoroo Potorous tridactylus trisulcatus	Gully line and patches of dense vegetation	Portland-Horsham FMA	Protect gully lines and patches of dense vegetation in areas where this species is known to occur and that are currently not protected through SPZ or SMZ .
Mallacoota Burrowing Crayfish Engaeus mallacoota	Known site	East Gippsland FMA	Apply a 100 m buffer from each bank for 1 km upstream and 1 km downstream of known Mallacoota Burrowing Crayfish sites. Avoid road construction and stream crossings within these stream buffers .
Malleefowl Leipoa ocellata	Active nesting mound	Portland-Horsham FMA	Apply a protection area of 3 ha and a management area of 250-300 m radius over each active nesting mound.

Species Name	Value	Applicable FMAs	Management Actions
Masked Owl Tyto novaehollandiae	Nesting and roosting site utilised recently and frequently or Old hollow-bearing tree	Central Highlands FMAs Midlands FMA	Apply a management area of 250 m radius over each nesting and roosting site utilised recently and frequently and located outside a Masked Owl Management Area, unless already protected. Within 250 m of the nesting or roosting site, protect habitat elements such as old hollow bearing trees that may be used by Masked Owls and their prey. Exclude timber harvesting operations, road construction and other activities likely to disturb breeding activity during the breeding season. Buffer nest trees by 100 m.
	Nesting and roosting site utilised recently and frequently or Habitat element	Gippsland FMAs North East FMAs Portland-Horsham FMA East Gippsland FMA	Apply a protection area of 3 ha and a management area of 250-300 m radius (or equivalent linear area) over each nesting and roosting site utilised recently and frequently and located outside a Masked Owl Management area, unless already protected. (In these cases, habitat for foraging is already provided in areas excluded from timber harvesting by general prescription including wildlife corridors, steep areas and unmerchantable areas and areas protected for other management purposes). Within 250-300 m of nesting or roosting site, protect habitat elements such as old hollow bearing trees that may be used by Masked Owls and their prey. Exclude timber harvesting operations, road construction and burning during the breeding season.
	Nesting and roosting site utilised recently and frequently or Habitat element	Otways FMA	Apply a protection area of 3 ha over each nesting and roosting site utilised recently and frequently and located outside a Masked Owl Management Area, unless already protected. (In these cases, habitat for foraging is already provided in areas excluded from timber harvesting operations by general prescription including wildlife corridors , steep areas and unmerchantable areas and areas protected for other management purposes). Within 250-300 m of nesting or roosting site, protect habitat elements such as old hollow bearing trees that may be used by Masked Owls and their prey.
Mount Stirling stonefly Thaumatoperla flaveola	Permanent stream	Gippsland FMAs	In known Mount Stirling stonefly habitat apply: -50 m buffers to permanent streams on the boundary of the logging coupe30 m buffers to permanent streams within the boundary of a logging coupe. Where the permanent stream commences within the coupe, apply the buffer for a distance of 30 m above the head of the permanent stream15 m filter strips to drainage lines. Where the drainage line commences within the coupe, apply the filter strip for a distance of 15 m above the head of the drainage line.

Species Name	Value	Applicable FMAs	Management Actions
Mountain Dragon Rankinia diemensis	Population	Midlands FMA	Apply a management area over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Mountain Galaxias Galaxias olidus	Permanent or temporary stream, or drainage line within a catchment containing Mountain Galaxias	Gippsland FMAs	Minimise stream crossings over permanent and temporary streams and drainage lines in catchments containing populations of Mountain Galaxias.
	Population	North East FMAs	Apply minimum stream buffer and filter strip widths specified in Table 10 for 1 km upstream of Mountain Galaxias populations or SMZ for Mountain Galaxias. Minimise stream crossings over waterways in catchments containing Mountain Galaxias.
	Population	East Gippsland FMA	Apply a protection area extending 100 m from each bank for 1 km upstream and 1 km downstream of populations. Avoid constructing new roads in the protection area . Note: The Secretary intends to review this strategy when 50 significant fish sites have been located.
Mountain Pygmy-possum Burramys parvus	Population	North East FMAs	Apply a management area over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Nankeen Night Heron Nycticorax caledonicus	Current roosting and breeding site	Mid-Murray FMA	Exclude activities likely to disturb breeding activity within 250 m radius of current roosting and breeding sites of colonially-nesting waterbirds during the breeding season.
Narracan Burrowing Crayfish Engaeus phyllocercus	Narracan Burrowing Crayfish site	Gippsland FMAs	Avoid road construction and stream crossings within stream buffers for 1 km upstream of Narracan Burrowing Crayfish sites. Design snig tracks to minimise alteration to drainage patterns in coupes adjacent Narracan Burrowing Crayfish sites on floodplain or seep zones. Avoid burning during late spring and early summer. Conduct burns when the moisture differential between Narracan Burrowing Crayfish habitat and the coupe is adequate to prevent the habitat from burning.

Species Name	Value	Applicable FMAs	Management Actions
Native fish	Major, permanent or temporary streams of the Aire River, Carlisle River and Gellibrand River	Otways FMA	For timber harvesting coupes adjacent Aire River, Carlisle River and Gellibrand River apply: - 100 m buffers to sections considered major streams; - 50 m buffers to sections considered permanent streams; and - 10 m buffers to sections considered temporary streams or greater if required to mitigate water quality risk.
New Holland Mouse Pseudomys novaehollandiae	Preferred habitat	Gippsland FMAs	Apply a protection area of 100 ha of preferred habitat including the detection site wherever possible.
Orbost Spiny Crayfish Euastacus diversus	Detection site	East Gippsland FMA	Apply a protection area extending 100m from each bank for 1 km upstream and 1 km downstream of detection sites. Avoid constructing new roads and stream crossings within the protection area . Manage nearby regeneration burns to ensure the protection area is not burnt.
			Note: The Secretary intends to review this strategy when 20 sites are established.
Otway Stonefly Eusthenia nothofagi	critical habitat	Otways FMA	In timber harvesting coupes within critical habitat of Otway Stonefly apply: - 100 m buffers to sections of the Aire River, Carlisle River, Gellibrand River considered major streams; - 50 m buffers to permanent streams; and - 10 m buffers to temporary streams or greater if required to mitigate water quality risk
Pale Sun Moth Synemon selene	Population	Portland-Horsham FMA	Apply a management area over populations. Detailed planning is required to ensure population is protected from timber harvesting.
Peregrine Falcon Falco peregrinus	Nesting site	East Gippsland FMA	Apply a management area of 250 m radius over each nesting site. Within 250 m of the nesting site, avoid timber harvesting, road construction and burning during breeding season. Apply a protection area of 100m radius around nesting site.
	Nest tree	Otways FMA	Buffer nest trees by 100 m, and avoid burning at all times.
Powerful Owl Ninox strenua	Nesting and roosting site	Central Highlands FMAs Midlands FMA Gippsland FMAs North East FMAs	Apply a protection area of 3 ha and a management area of 250-300 m radius (or equivalent linear area) around each nesting and roosting site, unless already protected.

Species Name	Value	Applicable FMAs	Management Actions
		Portland-Horsham FMA Bendigo FMA East Gippsland FMA Mid-Murray FMA	Outside of Powerful Owl Management Areas, habitat for foraging is provided in areas excluded from timber harvesting by general prescription including wildlife corridors , steep areas and unmerchantable areas and areas protected for other management purposes. Avoid burning operations within 250 m of known nest sites during the breeding season (1 May to 30 November).
			Within 250-300 m of nesting or roosting site, exclude timber harvesting operations, road construction and burning during the breeding season.
			Note: The Secretary intends to review zoning associated with individual nesting and roosting sites within the Central Highlands FMAs every 5 years. This review should prioritise the retention of zones for nesting and roosting sites utilised frequently within the previous 5 years.
	Nesting and roosting site	Otways FMA	Apply a protection area of 3 ha over each nesting and roosting site, unless already protected.
Regent Honeyeater Anthochaera phrygia	Detections outside of known sites	Midlands FMA	If Regent Honeyeaters are present in an area not listed as a regularly used site, exclude the area from timber harvesting operations until its significance for Regent Honeyeaters can be assessed by the Recovery Team and appropriate prescriptions devised.
	Regularly used sites (Regent Honeyeater)	Bendigo FMA Mid Murray FMA North East FMAs Portland-Horsham FMA	Apply a protection area extending 100 m from regularly used sites (Regent Honeyeater) . Surround the protection area with a management area of 150 m.
Rich reptile and amphibian sites	Site that is particularly rich in reptiles or amphibians	East Gippsland FMA	Apply a protection area of at least 10 ha at sites as being particularly rich in reptiles or amphibians
Royal Spoonbill Platalea regia	Current roosting and breeding site	Mid-Murray FMA	Exclude activities likely to disturb breeding activity within 250 m radius of current roosting and breeding sites of colonially-nesting waterbirds during the breeding season.
			Apply a management area over populations.
Samphire Skink Morethia adelaidensis	Population	Midlands FMA Portland Horsham FMA	Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
			Detailed planning is required to ensure population is protected from timber harvesting.

Species Name	Value	Applicable FMAs	Management Actions
Significant fish site	Site supporting 3 or more threatened fish species, or site supporting 6 or more native freshwater fish species	East Gippsland FMA	Apply a protection area extending 100 m from each bank for 1 km upstream and 1 km downstream of sites supporting 3 or more threatened fish species, or sites supporting 6 or more native freshwater fish species. Avoid constructing new roads in the protection area . Note: The Secretary intends to review this strategy when 50 significant fish sites have been located.
Small Brown Azure Butterfly Ogyris otanes	Host plant Berry broombush (Choretrum glomeratum)	Portland-Horsham FMA	Protect the host plant Berry broombush (<i>Choretrum glomeratum</i>) within 250m of any records of this species.
	Record	Central Highlands FMAs Gippsland FMAs North East FMAs Portland-Horsham FMA	Apply a management area of approximately 100 ha over records incorporating the detection site wherever possible.
Smoky Mouse Pseudomys fumeus			Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations or road construction commencement, incorporating any relevant information from studies of the species.
			Note: The Secretary intends to review this strategy when 10 Smoky Mouse SMZ are established or in light of further research regarding the conservation status of the species and its response to disturbance.
Sooty Owl Tyto tenebricosa	Nesting and roosting site utilised recently and frequently and located outside a Sooty Owl Management area, unless already protected	Central Highlands FMAs	Apply a management area of 250 m radius over each nesting and roosting site utilised recently and frequently and located outside a Sooty Owl Management area, unless already protected. In these cases, habitat for foraging is already provided in areas excluded from timber harvesting by general prescription including wildlife corridors, steep areas and unmerchantable areas and areas protected for other management purposes.
			Within 250 m of nesting or roosting site, exclude timber harvesting operations , road construction and other activities likely to disturb breeding activity during the breeding season. Buffer nest trees by 100 m.
	Nesting and roosting site utilised recently and frequently and located outside a Sooty Owl Management area, unless already protected	Gippsland FMAs North East FMAs East Gippsland FMA	Apply a protection area of 3 ha and a management area of 250-300 m radius (or equivalent linear area) over each nesting and roosting site utilised recently and frequently and located outside a Sooty Owl Management area, unless already protected. In these cases, habitat for foraging is already provided in areas excluded from timber harvesting by general prescription including wildlife corridors, steep areas and unmerchantable areas and areas protected for other management purposes. Within 250-300 m of nesting or roosting site, exclude
			vvitnin 250-300 m or nesting or roosting site, exclude

Species Name	Value	Applicable FMAs	Management Actions
			timber harvesting operations, road construction and burning during the breeding season.
South Gippsland Spiny Crayfish Euastacus neodiversus	Site	Gippsland FMAs	Develop management actions to protect South Gippsland Crayfish habitat at sites in State forest .
Southern Barred Frog Mixophyes balbus	Record	East Gippsland FMA	Where records of Southern Barred Frog are located on first-order streams or sites away from streams, protect approximately 50 ha (preferably the entire sub catchment unit) with a protection area . Where records of Southern Barred Frog are located on second or higher order streams, apply a protection area of 100 m width each side of the stream for 1 km upstream and downstream of the detection site.
			Avoid constructing new roads in the protection area.
			Note: The Secretary intends to review this strategy when an Action Statement is prepared.
Southern Bent- wing Bat Miniopteris schreibersii	Breeding and roosting cave, mine, or other over-wintering site	North East FMAs	Apply a protection area of 100 m radius over breeding and roosting caves and mines, and over-wintering sites.
Southern Brown Bandicoot Isoodon obesulus obesulus	Gully line and patches of dense vegetation	Portland-Horsham FMA	Protect gully lines and patches of dense vegetation in areas where this species is known to occur and that are currently not protected through SPZ or SMZ .
South-eastern Long-eared Bat Nyctophilus corbeni	Colony	Portland-Horsham FMA	Apply a 100 m buffer around colonies.
	Breeding and roosting site	Central Highlands FMAs	Apply and maintain a 100 m buffer around colonies.
Southern Myotis Myotis macropus	Breeding and roosting site	North East FMAs	Apply a protection area of 300 m radius over breeding and roosting sites. Note: The Secretary intends to review this strategy when 20 sites are established or when significant information has been obtained from further research.
	Breeding and roosting site	Gippsland FMAs	Apply a protection area of 100 m radius over breeding and roosting caves, trees and mines and over-wintering sites.
			_

Species Name	Value	Applicable FMAs	Management Actions
			Note: The Secretary intends to review this strategy when 20 sites are established or when significant information has been obtained from further research.
			Apply a management area of 200 m radius over breeding and roosting populations.
	Breeding and roosting site	Mid-Murray FMA	Within 200 m of breeding or roosting site, exclude all potentially disturbing activities during breeding season. Breeding season is 1 October to 31 March. Apply a 100m Buffer around breeding and roosting trees.
		East Gippsland	Apply a protection area of 100 m radius over breeding and roosting caves, trees and mines and over-wintering sites
			Apply a protection area of 500 ha and a management area of 1000 ha for all Quoll records in State forest throughout Victoria, up to a target of 50 sites.
Spot-tailed Quoll Dasyurus maculatus maculatus	Records or Preferred habitat feature or Detection site outside the SPZ within the SMZ, unless there are compelling reasons for excluding them	Central Highlands FMAs	Note: The location of the protection area and management area will be based on protecting preferred habitat features for Quolls. Protect den and latrine sites by a protection area of at least a 200 m radius. The protection area may include other detection sites, based on habitat quality and the proximity of existing protected habitat. Include detection sites outside the protection area within the management area, unless there are compelling reasons for excluding them (e.g. a record in a clearly unsuitable location for habitat protection, proximity of existing protected habitat etc.). Prioritise site protection for Quolls according to habitat quality, current reservation status of the site, linkage to other reserves and the presence of complementary values. Records within 2 km of each other are generally regarded as the same animal unless proved otherwise. Exclude the use of threatening poisons within 1 km of a record less than 5 years old.
			Note: The Secretary intends to periodically review the selection of Quoll records afforded SPZ and SMZ protection, to ensure that once targets are reached and as new records accrue or other information becomes available, the network of protected habitat in the FMA is optimal for Quoll conservation. Substitution of protected Quoll sites must consider the extent and quality of habitat and the currency, reliability and type of record.
	Records or	Gippsland FMAs North East FMAs	Apply a protection area of approximately 500 ha for each record. Include areas of undisturbed mature forest, riparian areas and rocky outcrops in the protection area where possible. In addition, apply a management area of approximately 1000 ha,

Species Name	Value	Applicable FMAs	Management Actions
	Preferred habitat feature		contiguous to the protection area or SPZ . Protect den and latrine sites by at least a 200 m radius protection area where they are not otherwise protected.
	Den and latrine site		Note: The objectives of the management area will be to maintain habitat for both arboreal and non-arboreal Spot-tailed Quoll prey, while providing continuing opportunities for timber harvesting. Harvesting within the management area or SMZ may comprise selective harvesting operations over the entire area, or carefully sited and scheduled seed tree or clearfall harvesting, ensuring, in either case that important prey habitat components such as hollow bearing trees are retained. Where clearfall or seed tree harvesting systems are proposed, harvesting operations may through time extend over the entire management area or SMZ. However, harvesting must be scheduled to ensure at least 500 ha of suitable prey habitat is available at any point in time (in addition to the protection area or SPZ). The adoption of harvesting strategies for management areas and SMZs should have regard to the assessed significance of the zone as Spot-tailed Quoll habitat, existing patterns of public land use and forest zoning, the silvicultural characteristics of forest types in the zone, the proportion of the zone which is unproductive for sawlog harvesting and the timber resource values of the productive areas.
			Note: The Secretary intends to review the selection of Quoll records afforded SPZ and SMZ protection when 10 zones have been established in State forest or when significant information has been obtained from further research. the purpose of the review is to ensure that once targets are reached and as new records accrue or other information becomes available, the network of protected habitat in the FMA is optimal for Quoll conservation. Substitution of protected Quoll sites must consider the extent and quality of habitat and the currency, reliability and type of record.
	Records		Apply a protection area of 500 ha and a management area of 1000 ha for all Quoll records in State forest , up to a target of 10 sites. The location of the protection area and management area will based on protecting preferred habitat features for Quolls. Protect den and latrine sites by a protection area of at least a 200 m radius.
	Preferred habitat feature Den and latrine site	Portland-Horsham FMA	Note: The protection area may include other detection sites, based on habitat quality and the proximity of existing protected habitat. Include detection sites outside the protection area within the management area , unless there are compelling reasons for excluding them (e.g. a record in a clearly unsuitable location for habitat protection, proximity of existing protected habitat
			etc.). Prioritise site protection for Quolls according to habitat quality, current reservation status of the site, linkage to other reserves and the presence of

Species Name	Value	Applicable FMAs	Management Actions
			complementary values. Records within 2 km of each other are generally regarded as the same animal unless proved otherwise. Exclude the use of poisons such as 1080 within 2 km of Quoll records.
			Apply a protection area of 500 ha and a management area of 1000 ha for all Quoll records in State forest up to a target number of sites (to be determined). The location of the protection area and management area will be based on protecting preferred habitat features for Quolls. The protection area will include known den and latrine sites (protected by at least a 200 m radius), and may include other detection sites, based on habitat quality and the proximity of existing protected habitat.
	Preferred habitat feature or Detection site	East Gippsland FMA	Note: Detection sites not included in the protection area will generally be included within the management area, unless there are compelling reasons for excluding them (e.g. a record in a clearly unsuitable location for habitat protection, proximity of existing protected habitat etc.). Site protection for Quolls will be prioritised according to habitat quality, current reservation status of the site, linkage to other reserves and the presence of complementary values. Records within 2 km of each other will be generally regarded as the same animal unless proved otherwise.

Note: The Secretary intends to periodically review the selection of Quoll records afforded SPZ and SMZ protection, to ensure that once targets are reached and

Species Name	Value	Applicable FMAs	Management Actions
			Apply a protection area of 500 ha and a management area of 1000 ha for all Quoll records in State forest throughout Victoria, up to a target of 10 sites.
	Preferred habitat feature or Detection site	Midlands FMA	Note: The location of the protection area and management area will based on protecting preferred habitat features for Quolls. Protect den and latrine sites with a protection area of at least a 200 m radius. The protection area may include other detection sites, based on habitat quality and the proximity of existing protected habitat. Include detection sites outside the protection area within the management area, unless there are compelling reasons for excluding them (e.g. a record in a clearly unsuitable location for habitat protection, proximity of existing protected habitat etc.). Prioritise site protection for Quolls according to habitat quality, current reservation status of the site, linkage to other reserves and the presence of complementary values. Records within 2 km of each other are generally regarded as the same animal unless proved otherwise. Exclude the use of poisons such as 1080 within 2 km of Quoll records. Note: The Secretary intends to periodically review the selection of Quoll records afforded SPZ and SMZ protection, to ensure that once targets are reached and as new records accrue or other information becomes available, the network of protected habitat in the FMA is optimal for Quoll conservation. Substitution of protected Quoll sites must consider the extent and quality of habitat and the currency, reliability and type of record.
	Preferred habitat feature or Detection site	Otways FMA	Apply a protection area of approximately 500 ha for each record. The location of the protection area will be based on protecting preferred habitat features for Quolls. The protection area will include known den and latrine sites (protected by at least a 200 m radius), and may include other detection sites, based on habitat quality and the proximity of existing protected habitat. Note: Site protection for Quolls will be prioritised according to habitat quality, current reservation status of the site, linkage to other reserves and the presence of complementary values. Records within 2 km of each other will be generally regarded as the same animal unless proved otherwise. Note: The Secretary intends to periodically review the selection of Quoll records afforded SPZ and SMZ protection, to ensure that once targets are reached and as new records accrue or other information becomes available, the network of protected habitat in the FMA is optimal for Quoll conservation. Substitution of protected Quoll sites must consider the extent and quality of habitat and the currency, reliability and type of record.

Species Name	Value	Applicable FMAs	Management Actions
Spotted Galaxis Galaxius truttaceaus	Population	Midlands FMA	Apply a management area for populations.
Spotted Tree Frog Litoria spenceri	Locations where frogs have been recorded and where suitable habitat has been mapped	Central Highlands FMAs	Until the species critical habitat is known, apply a protection area of 300 m radius over locations where frogs have been recorded and where suitable habitat has been mapped (collectively referred to as 'stream habitat'). Also exclude new road construction and other potentially threatening activities (such as prescribed fire) from this area. For at least 1km upstream from and potential Spotted Tree Frog locations: - apply buffer and filter widths set out in Appendix 1 Table 10 to timber harvesting operations; and - do not construct new roads or stream crossings. In the catchment upstream of a Spotted Tree Frog management zone: - ensure any new road (and the fill slope toe of any new road) is located at least 50 m from any stream, unless site specific sediment management operations are put in place to prevent sediments entering perennial and ephemeral streams in the management zone - minimise construction of new roads and stream crossings, and ensure any constructed comply with the following prescriptions extracted from O'Shaughnessy and Associates (1997): • Ensure that creek culverts and bridges are built to be effective with 1:50 year flows. • Construct rock armoured overflow sections to allow overtopping without road washouts. • Install culverts of a minimum dimension of 375 mm along roads.
			 500 mm either side of stream crossings and construct crossings to ensure that road drainage at the crossing does not directly enter the stream. Ensure that all culvert outlets have structures in place to spread and disperse culvert discharge using materials such as rocks, devises constructed from durable timber and logs.

durable timber and logs.

Use excavators to prepare stream crossings for culvert placement. Ensure that material is pulled back on either side and not across the crossing. Ensure that fill placement is accurate following culvert.	Species Name	Value	Applicable FMAs	Management Actions
• Ensure that fill placement is accurate following culturat				culvert placement. Ensure that material is pulled back
installation.				• Ensure that fill placement is accurate following culvert installation.
 Use techniques to ensure silt runoff from surface flows, embankments and fills into drainage lines and or creeks is minimised. 				flows, embankments and fills into drainage lines and
 When removing stream crossings associated with temporary roads: (a) extensively drain the road formation using substantial banks: and (b) remove the road bench where it connects to the permanent road network. 				temporary roads: (a) extensively drain the road formation using substantial banks: and (b) remove the road bench where it connects to the permanent road
- schedule timber harvesting operations , roading and burning to minimise the area disturbed at any one time				and burning to minimise the area disturbed at any one
Until the species critical habitat is known, apply a protection area of 300 m radius over all frog sites and mapped habitat areas. Apply a management area of a further 700m either side of the protection area and 1 km upstream or to the ridge top (if the stream is less than 1 km long) along all streams that flow into the frog habitat. Exclude timber harvesting operations, new road construction and other potentially threatening activities (such as prescribed fire) from the protection area.				protection area of 300 m radius over all frog sites and mapped habitat areas. Apply a management area of a further 700m either side of the protection area and 1 km upstream or to the ridge top (if the stream is less than 1 km long) along all streams that flow into the frog habitat. Exclude timber harvesting operations, new road construction and other potentially threatening activities (such as prescribed fire) from the protection
Within the management area or SMZ:				Within the management area or SMZ:
Locations where frogs have been recorded and where suitable habitat has been mapped Gippsland FMAs North East FMAs Gippsland FMAs North East FMAs - do not construct new stream crossings - ensure any new road (and the fill slope toe of any new road) is located at least 50 m from any stream, unless site specific sediment management operations are put in place to prevent sediments entering perennial and ephemeral streams in the management zone - apply buffer and filter widths set out in Table 10 give particular attention to suspension of logging and snig track drainage during wet weather - excavate 50 cm deep pits below the original track surface and immediately upslope of bars constructed to drain snig tracks and other tracks - ensure landings are constructed, or are drained to dispose of discharge at least 50 m from the upslope edge of buffers and filters.		have been recorded and where suitable habitat has been	• •	 ensure any new road (and the fill slope toe of any new road) is located at least 50 m from any stream, unless site specific sediment management operations are put in place to prevent sediments entering perennial and ephemeral streams in the management zone apply buffer and filter widths set out in Table 10. give particular attention to suspension of logging and snig track drainage during wet weather excavate 50 cm deep pits below the original track surface and immediately upslope of bars constructed to drain snig tracks and other tracks ensure landings are constructed, or are drained to dispose of discharge at least 50 m from the upslope

Species Name	Value	Applicable FMAs	Management Actions
	Nest tree used within the last 5 years	Central Highlands FMAs North East FMAs Portland-Horsham FMA	Apply a 100 m buffer around nest trees used within the last 5 years. Exclude timber harvesting operations within 250 m of nest trees during breeding season.
Square-tailed Kite			Apply a management area of 250 m radius over nesting sites.
Lophoictinia isura	Nesting site	Midlands FMA Gippsland FMAs Bendigo FMA East Gippsland FMA	Within 250 m of nesting sites, avoid timber harvesting, road construction and burning during breeding season. Apply a protection area of 100m radius around nesting sites.
			Note: Within the Midlands FMA , the Secretary intends to review this strategy when 10 sites have been established in State forest .
	Population		Apply a management area of approximately 500 ha of suitable habitat over populations.
Squirrel Glider Petaurus norfolcensis	Key habitat element such as nesting hollows, large trees and suitable understorey species	North East FMAs Portland-Horsham FMA Midlands FMA	Within the management area or SMZ, implement prescriptions that address maintenance of key habitat elements such as nesting hollows, large trees and suitable understorey species (in particular Silver Wattle Acacia dealbata). Exclude timber harvesting operations from a core area of approximately 100 ha. Note: The Secretary intends to review this strategy when 10 sites have been established in State forest.
	Population	Bendigo FMA	Apply a management area over populations. Within the management area or SMZ, implement prescriptions that address maintenance of key habitat elements such as nesting hollows, large trees and suitable understorey species.
Strzelecki Burrowing Crayfish Engaeus rostrogaleatus	Sites	Gippsland FMAs	Develop management actions to protect Strzelecki Burrowing Crayfish habitat at sites in State forest .
			Apply a management area of 250 m radius over nest trees.
Superb Parrot Polytelis swainsonii	Nest tree	Mid-Murray FMA	Within 250m of the nest tree, exclude all potentially disturbing activities during breeding season. Breeding season is 1 September to 31 December. Apply a protection area of 100m radius around nest trees.
			Note: The Secretary intends to review this strategy statewide every 10 years from 2003, or if the total area of buffer zones in Victorian State forests exceeds 200 ha, whichever occurs first.

Species Name	Value	Applicable FMAs	Management Actions
Swamp Skink Lissolepis coventryi	Records	Portland-Horsham FMA	Minimise disturbance to damp areas and gully lines. Maintain dense vegetation within 250 m of records of this species.
Swift Power	Vicinity of Swift Parrot flocks	Midlands FMA Portland-Horsham FMA	Avoid disturbance of Swift Parrot flocks by postponing timber harvesting operations that may affect them in their vicinity.
Swift Parrot Lathamus discolor			Apply a management area for resident populations.
uiscoloi	Resident population	North East FMAs	Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations or prescribed burning activities.
Trout Cod Maccullochella macquariensis	Viable Trout Cod population	North East FMAs Mid-Murray FMA	Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately managed and protected when timber harvesting operations are planned in rivers and catchments upstream of viable Trout Cod populations.
Turquoise Parrot Neophema Pulchella	Stump identified as nest site	North East FMAs	Protect stumps identified as nest sites from timber harvesting operations.
Variegated Pygmy Perch Nannoperca	Population	Portland-Horsham FMA	Apply a protection area extending 40 m from each bank for 40 m upstream and 40 m downstream of populations.
variegata			Disturbance that could impact on water quality must be avoided within a protection area .
Warragul			Apply a management area over populations.
Burrowing Crayfish Engaeus sternalis	Population	Gippsland FMAs	Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Wedge-tailed Eagle Aquila audax	Nesting site	Midlands FMA	Apply a management area of 250 m radius over nesting sites. Within the management area or SMZ, avoid timber harvesting, road construction and burning activities during breeding season. Breeding season is 1 July to 30 November. Within 100m of nest trees, exclude timber harvesting operations, road construction and burning at all times.
			Note: The Secretary intends to review this strategy when 10 sites have been established in State forest .

Species Name	Value	Applicable FMAs	Management Actions
			Apply a management area of 500 m radius over nesting sites.
	Nesting site	Gippsland FMAs	Within 500m of nesting sites, avoid timber harvesting, road construction and burning during breeding season. Apply a protection area of 100m radius around nesting sites.
			Apply a management area over resident populations.
White-bellied	Resident population	North East FMAs	Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Sea-eagle Haliaeetus Ieucogaster		Portland-Horsham	Apply a management area of 250 m radius over each nesting site.
	Nesting site	FMA East Gippsland FMA	Within 250m of nesting site, avoid timber harvesting, road construction and burning during breeding season. Apply a protection area of 100m radius around nesting sites.
	_		Apply a management area of 250 m radius over nesting populations.
	Nesting population Mid-Murray FM		Within 20m of nesting population, avoid timber harvesting, road construction and burning during breeding season. Breeding season is 1 May to 31 December. Apply a protection area of 100 m radius around nesting sites.
White-footed Dunnart Sminthopsis leucopus	Site	Portland-Horsham FMA	Apply a protection area of approximately 20 ha of suitable habitat over sites.
	Site	North East FMAs	Apply a protection area of approximately 50 ha over each site, where practicable incorporating the detection site.
Woodland Blind Snake Anilios proximus	NOTH Last Pivir		Note: The Secretary intends to review the strategy when 20 SPZ are established.
	Population Mid-Murray FMA		Apply a management area of approximately 50 ha over populations, where practicable incorporating the detection site.
			Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .

Species Name	Value	Applicable FMAs	Management Actions
	Population	Midlands FMA	Apply a management area over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
	Den tree and associated flight paths, and feed trees	Portland-Horsham FMA	Retain all Yellow-Bellied Glider den trees and associated flight paths, and feed trees. (Yellow-Bellied Glider feed trees are recognised by "V – notch" incisions into the sapwood and are found in a range of species including <i>Eucalyptus viminalis</i> , <i>E. ovata</i> , <i>E. baxteri</i> , <i>E. obliqua and E. willisii</i> .)
Yellow-bellied Glider Petaurus australis	Relative abundance (More than 5 per Spotlight Kilometre)	East Gippsland FMA Otways FMA	Apply a protection area of approximately 100 ha of suitable habitat where records report a relative abundance of more than 5 individuals per spotlight kilometer (equivalent to more than 0.2 individuals per hectare or more than 7 individuals per hour of spotlighting), or where substantial populations are located in isolated or unusual habitat. Note: Assumed rate of spotlighting per kilometer is 10mins per 100m and visible range either side of transect is 150m, equating to assumed minimum survey
	Roost site with patterns of regular seasonal use	Gippsland FMAs	Apply a protection area of 100 m radius over roost sites with patterns of regular seasonal use. Establish a further management area of 100 m radius around the protection area for sites of active colonies.
Yellow-bellied Sheathtail Bat Saccolaimus flaviventris	Sites of active colonies		Schedule timber harvesting, roading and fuel reduction burning operations within the management area to minimise disturbance to colonies.
	Older forest over sites where more than 10 individuals are captured in one trapping session	East Gippsland FMA	Apply a protection area of approximately 100 ha of older forest over sites where more than 10 individuals are captured in one trapping session. Note: The Secretary intends to review this strategy when 20 sites are established.

Table 14 Rare or threatened flora prescriptions

Species Name	Value	Applicable FMAs	Management Actions
Alpine Bush-pea Pultenaea fasciculata	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Alpine Buttons Leptorhynchos squamatus subsp. Alpinus	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Alpine Colobanth Colobanthus affinis	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Alpine Crane's-bill Geranium brevicaule	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Alpine Fen-sedge Carex hypandra	Population	North East FMAs	Apply a protection area of 200 m radius over each population.
Alpine Finger-fern Notogrammitis crassior	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Alpine Groundsel Senecio pectitus var. major	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Alpine Marianth Rhytidosporum inconspicuum	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Value	Applicable FMAs	Management Actions
Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Population	Gippsland FMAs North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Population	Otways FMA	Manage occurrences in consultation with the Department unless already protected. Implement hygiene precautions to control the potential introduction and / or spread of <i>Phytophthora cinnamomic</i> .
Population	Gippsland FMAs	Apply a protection area over each population.
Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection
	Population Population Population Population Population Population Population Population	Population North East FMAs Population North East FMAs Population North East FMAs Population North East FMAs Population Gippsland FMAs North East FMAs Population Otways FMA Population Gippsland FMAs North East FMAs

Species Name	Value	Applicable FMAs	Management Actions
			and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Ausfeld's Wattle Acacia ausfeldii	Population	Bendigo FMA	Minimise disturbance from road management activities.
Austral Crane's-bill Geranium solanderi var. solanderi s.s.	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Austral Moonwort Botrychium australe	Population	North East FMAs East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
	Population	Gippsland FMAs North East FMAs	Apply a protection area over each population.
Austral Toad-flax Thesium australe	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Austral Trefoil Lotus australis var. australis	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Australian Anchor Plant Discaria pubescens	Population	North East FMA East Gippsland FMA	Apply a management area of 200m over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
	Population	Gippsland	Apply a protection area over each population.
Bantam Bush-pea Pultenaea parrisiae	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Basalt Peppercress Lepidium hyssopifolium s.s.	Population	Midlands FMA	Apply a protection area over each population. Conduct pre-harvest surveys for the species in the vicinity of Bolworrah and Trentham.
Bassian Pomaderris Pomaderris oraria subsp. oraria	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Baw Baw Sally Eucalyptus pauciflora subsp. acerina	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Bead Glasswort Tecticornia flabelliformis	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Beechworth Silver Stringybark Eucalyptus cinerea subsp. Victoriensis	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Ben Major Grevillea Grevillea floripendula	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.
Bent-leaf Wattle Acacia flexifolia	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Betka Bottlebrush Callistemon kenmorrisonii	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Binung Christella dentata	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Birch Pomaderris Pomaderris betuli subsp. Betulina	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.

Species Name	Value	Applicable FMAs	Management Actions
Black Oliveberry Elaeocarpus holopetalus	Population	Portland- Horsham FMA	Manage populations in consultation with the Department unless already protected.
Black Stem Adiantum formosum	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Blackfellow's Hemp Androcalva rossii	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Blotched Hyacinth- orchid Dipodium variegatum	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Blue-leaf Tussock- grass Poa sieberiana var. cyanophylla	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Blue-tongue Greenhood Pterostylis oreophila	Population	Gippsland FMAs North East FMAs	Apply a protection area of 200 m radius over each population.
Blunt Sandalwood Santalum obtusifolium	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Blunt-leaf Pomaderris Pomaderris helianthemifolia	Population	Gippsland FMAs North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Bog Saw-sedge Gahnia subaequiglumis	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Bogong Daisy-bush Olearia frostii	Population	North East FMAs	Apply a protection area of 200 m radius over each population.
Bonnet Orchid Cryptostylis erecta	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Branching Raspwort Gonocarpus micranthus subsp. ramosissimus	Population	North East FMAs	Protect populations from disturbance where possible.
			Apply a management area over populations.
Bristly Greenhood Pterostylis setifera	Population	Bendigo FMA	Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Bristly Helmet-orchid Corybas hispidus	Population	Gippsland FMAs North East FMAs	Apply a protection area of 200 m radius over each population.
Brittle Bladder-fern Cystopteris tasmanica	Population	North East FMAs	Apply a protection area of 200 m radius over each population.
Broad-leaf Hop-bush Dodonaea rhombifolia	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
			Apply a management area over populations.
Broad-lip Diuris Diuris X palachila	Population	Bendigo FMA	Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Broad-lip Leek-orchid Prasophyllum patens s.l.	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Brooker's Gum Eucalyptus brookeriana	Occurence	Otways FMA	Manage occurrences in consultation with the Department unless already protected. Where possible, reseed or replant all logging coupes that contained E. brookeriana with seed collected from the coupe to ensure reestablishment of this species on the site in a similar proportion to that found prior to harvesting as shown on the coupe plan.
Broom Bitter-pea Daviesia genistifolia s.s.	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.
Buff Hazelwood Symplocos thwaitesii	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Buffalo Sallee Eucalyptus mitchelliana	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Buloke Mistletoe Amyema linophylla subsp. orientale	Population	Mid-Murray FMA	Apply a management area of 100 m radius over populations that are not already protected. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Buloke Allocasuarina luehmannii	Population	Bendigo FMA	Minimise disturbance from road management activities.
Button Rush	Population	Mid-Murray FMA	Apply a management area of 100 m radius over populations that are not already protected. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Lipocarpha microcephala	Population	North East FMAs	Apply a management area of 200 m radius over populations that are not already protected. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Buxton Gum Eucalyptus crenulata	Population	Central Highlands FMAs	Protect populations from disturbance where possible.
Cabbage Fan-palm Livistona australis	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Candy Spider-orchid Caladenia versicolor	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Cane Spear-grass Austrostipa breviglumis	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.
- Tuoti ootipa Brovigianno	Population	Bendigo FMA	Minimise disturbance from road management activities.
Castlemaine Spider- orchid Caladenia clavescens	Population	Bendigo FMA	Apply a management area over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Catkin Wattle Acacia dallachiana	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Chinese Lespedeza Lespedeza juncea subsp. sericea	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Cliff Cudweed Euchiton umbricola	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
	Population	Midlands FMA	Apply a protection area over each population based on field inspection. Review this strategy when 10 sites and 1000 individuals are protected in State forest . Note: The Secretary intedns to review this
			strategy when 10 sites and 1000 individuals are protected in State forest
Clover Glycine Glycine latrobeana	Population	Gippsland FMAs North East FMAs	· · · · · · · · · · · · · · · · · · ·
		Portland-	Apply a management area over populations based on field inspection.
	Population	Horsham FMA Bendigo FMA	Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Coast Bitter-bush Adriana quadripartita (pubescent form)	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Coast Dandelion Taraxacum cygnorum	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Coast Grey-box Eucalyptus bosistoana	Coast Grey Box stand	Gippsland FMAs	Exclude timber harvesting operations from Coast Grey Box stands , except for when it is undertaken to facilitate propagation or regeneration of this species.
Coast Ixodia Ixodia achillaeoides subsp. arenicola	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Coastal Greenhood Pterostylis alveata	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Cobberas Grevillea Grevillea brevifolia	Population	Gippsland FMAs North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is

Species Name	Value	Applicable FMAs	Management Actions
			adequately protected during timber harvesting operations.
Cobra Greenhood Pterostylis grandiflora	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Colourful Spider- orchid Caladenia colorata	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Colquhoun Grevillea Grevillea celata	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Comb Wheat-grass Australopyrum retrofractum	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Common Joyweed Alternanthera nodiflora	Population	Mid-Murray FMA	Avoid disturbance to populations.
Common Spider-orchid Caladenia patersonii s.l.	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.
Common Spleenwort Asplenium trichomanes	Population	Gippsland FMAs North East FMAs	· · · · · · · · · · · · · · · · · · ·
Concave Pomaderris Pomaderris subplicata	Population	Gippsland FMAs North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
			Apply a management area of 200 m radius over populations.
Convex Pomaderris Pomaderris subcapitata	Population	Gippsland FMAs North East FMAs	Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Creeping Grevillea Grevillea repens	Population	Central Highlands FMAs	Consult with a botanist to determine if management actions are required to protect occurrences in the vicinity of prescribed burning activities.
-	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.
Crested Hair-grass Koeleria macrantha	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Crested Water-milfoil Myriophyllum lophatum	Population	North East FMAs	Protect populations from disturbance where possible.
Crimson Grevillea Grevillea polybractea	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Crimson Spider Orchid Caladenia concolor	Population	Central Highlands FMAs	Protect populations from disturbance where possible. Consult with a botanist to determine if management actions are required to protect occurrences in the vicinity of prescribed burning activities.
	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Cryptic Heath Epacris celata	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Cupped Bush-pea Pultenaea vrolandii	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the

Species Name	Value	Applicable FMAs	Management Actions
			Department to ensure the species is adequately protected during timber harvesting operations.
Curly Sedge Carex tasmanica	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Currawang Acacia doratoxylon	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Dark Wire-grass Aristida calycina var. calycina	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Deane's wattle Acacia deanei subsp. deanei	Population	North East FMAs	Apply a protection area of 200 m radius over each population.
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Delicate Bush-pea Pultenaea tenella	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Delicate Cranesbill Geranium sp. 6	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Delicate Love-grass Eragrostis tenellula	Population	Mid-Murray FMA	Apply a management area of 100 m radius over populations that are not already protected. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Delicate New Holland Daisy Vittadinia tenuissima	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is

Species Name	Value	Applicable FMAs	Management Actions
			adequately protected during timber harvesting operations.
Dense Leek-orchid Prasophyllum spicatum	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Dense Mint-bush Prostanthera decussata	Population	Gippsland FMAs North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Dergholm Guinea- flower Hibbertia humifusa subsp. debilis	Population	Portland- Horsham FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Devious Bent-grass Deyeuxia decipiens	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Dookie Daisy Brachyscome gracilis subsp. gracilis	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Downy Star-Bush Asterolasia phebalioides	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Dusky Bush-pea Pultenaea polifolia	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Dwarf Buttercup Ranunculus millanii	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Dwarf Milkwort Polygala japonica	Population	Gippsland FMAs North East FMAs East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Dwarf Swainson-pea Swainsona phacoides	Population	Mid-Murray FMA	Apply a management area of 100 m radius over populations that are not already protected. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Eastern Bitter-bush Adriana tomentosa var. tomentosa (glabrous and pubescent form)	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Eastern Pomaderris Pomaderris discolor	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Eichler's Buttercup Ranunculus eichlerianus	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Elegant Spider-orchid Caladenia formosa	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Elusive Cress Irenepharsus magicus	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Emerald-lip Greenhood Pterostylis smaragdyna	Population	Bendigo FMA	Apply a management area over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .

Species Name	Value	Applicable FMAs	Management Actions
Enfield Grevillea Grevillea bedgoodiana	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.
Erect Midge-orchid Corunastylis arrecta	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Errinundra Pepper Tasmannia xerophila subsp. robusta	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Fairy Bluebell Wahlenbergia densifolia	Population	Gippsland FMAs	Apply a protection area over populations.
	Site	Central Highlands FMAs	Apply a 20 m buffer around sites. On advice from the Department , buffers may not be required if locally abundant in nearby parks and reserves or SPZ .
Fairy Lanterns Thismia rodwayi	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Fairy Orchid Caladenia alata	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Ferny Small-flower Buttercup Ranunculus pumilio var. politus	Population	North East FMAs	Protect populations from disturbance where possible.
Filmy Maidenhair Adiantum diaphanum	Population	Gippsland FMAs	Apply a protection area over each population.
Fine Bent-grass Deyeuxia boormanii	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Fir Clubmoss Huperzia australiana	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is

Species Name	Value	Applicable FMAs	Management Actions
			adequately protected during timber harvesting operations.
Fisch's Greenhood Pterostylis fischii	Population	Gippsland FMAs North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Flat-leaf Bush-pea Pultenaea platyphylla	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Floating Bur-reed Sparganium subglobosum	Population	North East FMAs	Protect populations from disturbance where possible.
Floodplain Rustyhood Pterostylis cheraphila	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Fog Club-sedge Isolepis montivaga	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Forde Poa Poa fordeana	Population	Mid-Murray FMA	Avoid disturbance to populations.
Forest Weft-moss Thuidium laeviusculum s.s.	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Forrester's Bottlebrush Callistemon forresterae	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Fragrant Leek-orchid Prasophyllum suaveolens	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Fragrant Saltbush Rhagodia parabolica	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.

Species Name	Value	Applicable FMAs	Management Actions
Fringed Rice-flower Pimelea ligustrina subsp. ciliata	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Genoa River Correa Correa lawrencea var. genoensis	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Gippsland Hemp Bush Gynatrix macrophylla	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Gippsland Lakes Peppermint Eucalyptus arenicola	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Glandular Early Nancy Wurmbea biglandulosa subsp. biglandulosa	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Globe-hood Sun-orchid Thelymitra X chasmogama	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Golden Cowslips Diuris behrii	Population	Bendigo FMA	Apply a management area over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Golden Grevillea Grevillea chrysophaea	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.

Species Name	Value	Applicable FMAs	Management Actions
Golden Moths Diuris lanceolata s.l.	Population	Gippsland FMAs	Apply a protection area over each population.
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Golden Pomaderris Pomaderris aurea	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Goldfield Boronia Boronia anemonifolia subsp. Aurifodina	Population	Bendigo FMA	Minimise disturbance from road management activities.
Gorae Leek-orchid Prasophyllum diversiflorum	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Graceful Sun-orchid Thelymitra simulata	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Grampians Bitter-pea Daviesia laevis	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Grampians Pincushion-lily Borya mirabilis	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Grampians Rice-flower Pimelea pagophila	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Grampians Zieria Zieria oreocena	Population	Otways FMA	Manage occurrences in consultation with the Department unless already protected.
Granite Greenhood Pterostylis tunstallii	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Green Leek-orchid Prasophyllum lindleyanum	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is

Species Name	Value	Applicable FMAs	Management Actions
			adequately protected during timber harvesting operations.
Green Wattle Acacia irrorata subsp. irrorata	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Greenish-flower Vanilla-lily Arthropodium sp. 2 (greenish flowers)	Population	North East FMAs	Protect populations from disturbance where possible.
Green-striped Greenhood Pterostylis chlorogramma	Population	Gippsland FMAs Portland- Horsham FMA	Apply a protection area over each population.
Green-Top Sedge Carex chlorantha	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Grey Beard-heath Leucopogon attenuatus	Population	Gippsland FMAs North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Grey Rice-flower Pimelea treyvaudii	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Grey Scentbark Eucalyptus ignorabilis s.s.	Population	Gippsland FMAs	Apply a protection area of 200 m radius around each population.
Gully Grevillea Grevillea barklyana	Mature individual	Central Highlands FMAs	Protect mature individuals from disturbance where possible.
Gunn's Alpine Buttercup Ranunculus gunnianus	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Hairy Eyebright Euphrasia lasianthera	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Hairy Hop-bush Dodonaea boroniifolia	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Hairy-pod Wattle Acacia glandulicarpa	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Hard-head Bush-pea Pultenaea capitellata	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Heath Platysace Platysace ericoides	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Highland Bush-pea Pultenaea williamsonii	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Honey Hood-orchid Caladenia hildae	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Jerry-jerry Ammannia multiflora	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Jumping-jack Wattle Acacia enterocarpa	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.

Species Name	Value	Applicable FMAs	Management Actions
Kerrawang Commersonia dasyphylla	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
King Greenhood Pterostylis baptistii	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Kydra Dampiera Dampiera fusca	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Lady's Mantle Alchemilla xanthochlora	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
	Population	Gippsland FMAs North East FMAs	Apply a protection area of 200 m radius over each population.
Lanky Buttons Leptorhynchos elongatus	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Large Rustyhood Pterostylis maxima	Population	Bendigo FMA	Apply a management area over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Large Tick-trefoil Desmodium brachypodum	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Large Velvet Wallaby- grass Rytidosperma pilosum var. paleaceum	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Large-leaf Ray-flower Cyphanthera anthocercidea	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Leafless Pink-bells Tetratheca subaphylla	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Leafless Tongue- orchid Cryptostylis hunteriana	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
	Population	Gippsland FMAs	Apply a protection area over each population.
Leafy Greenhood Pterostylis cucullata	Population	North East FMAs Portland- Horsham FMA East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the population is adequately protected during timber harvesting operations.
Leafy Templetonia Templetonia stenophylla	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.
Leafy Wallaby-grass Rytidosperma bipartitum s.s.	Population	Bendigo FMA	Minimise disturbance from road management activities.
Liles Down	Population	Gippsland FMAs	Apply a protection area over each population.
Lilac Berry Trochocarpa clarkei	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is

Species Name	Value	Applicable FMAs	Management Actions
			adequately protected during timber harvesting operations.
Lilac Bitter-cress Cardamine lilacina s.s.	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Limestone Blue Wattle Acacia caerulescens	Population	Gippsland FMAs East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Limestone Pomaderris Pomaderris oraria subsp. calcicola	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Limestone Spider- orchid Caladenia calcicola	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Little Kooka Wattle Acacia nanopravissima ms.	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Long Rope-rush Calorophus elongatus	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Long-flower Beard- heath Leucopogon juniperinus	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
-	Population	North East FMAs	Protect populations from disturbance where possible.
Long-tail Greenhood Pterostylis woollsii	Population	Bendigo FMA	Apply a management area over populations. Conduct a site inspection and detailed planning in consultation with the Department

Species Name	Value	Applicable FMAs	Management Actions
			to ensure the species is adequately protected during timber harvesting operations .
Long-tongue Summer Greenhood Pterostylis aestiva	Population	Gippsland FMAs North East FMAs	
Low Bush-pea Pultenaea subspicata	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Lowan Phebalium Phebalium lowanense	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
			Apply a management area over populations.
Lowly Greenhood Pterostylis despectans	Population	Bendigo FMA	Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations . Within the management area , timber harvesting activities are limited to thinning operations conducted in accordance with plans developed for the specific purpose of aiding the survival of Lowly Greenhood.
Maiden's Wattle Acacia maidenii	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Mallee Ash Eucalyptus kybeanensis	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Maroon Leek-orchid Prasophyllum frenchii	Population	Portland- Horsham FMA Otways FMA	Manage occurrences in consultation with the Department unless already protected.

Species Name	Value	Applicable FMAs	Management Actions
	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Marsh Daisy Brachyscome radicans	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Marsh Greenhood Pterostylis uliginosa	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Marsh Leek-orchid Prasophyllum niphopedium	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Marsh Sun-orchid Thelymitra longiloba	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Mat Cudweed Euchiton traversii	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Matted Flax-lily Dianella amoena	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Matted Parrot-pea Dillwynia prostrata	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
McIvor Spider-orchid Caladenia audasii	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Mellblom's Spider- orchid Caladenia hastata	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Metallic Sun Orchid	Population	Gippsland FMAs	Apply a protection area over each population.
Thelymitra epipactoides	Occurrence	Portland- Horsham FMA Otways FMA	Manage occurrences in consultation with the Department unless already protected.
Monkey Mint-bush Prostanthera walteri	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Mossy Knawel Scleranthus singuliflorus	Population	Gippsland FMAs North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Mount Cole Grevillea Grevillea montis-cole	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.
Mountain Aciphyll Aciphylla simplicifolia	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Mountain Dandelion Taraxacum aristum	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Mountain Kangaroo Apple Solanum linearifolium	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Mountain Leafless	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Bossiaea bracteosa	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Mountain Leek-orchid Prasophyllum aff. montanum B	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Mountain Mat-rush Lomandra oreophila	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Mountain Swainson- pea Swainsona recta	Population	North East FMAs	Apply a protection area of 200 m radius over each population.
Mountain Wheat-grass Australopyrum velutinum	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Mueller Daisy Brachyscome muelleroides	Population	Mid-Murray FMA	Apply a management area of 100 m radius over populations that are not already protected. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Mueller's Bent Agrostis muelleriana	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Mugga Eucalyptus sideroxylon subsp. sideroxylon	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Naked Sun-orchid Thelymitra circumsepta	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Narrow Goodenia Goodenia macbarronii	Population	North East FMAs	Apply a protection area over each population.
Narrow-leaf Star-hair Astrotricha linearis	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Native Quince Alectryon subcinereus	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Native wintercress Barbarea grayi	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Netted Daisy-bush Olearia speciosa	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Omeo Gum Eucalyptus neglecta	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Orange-blossom Orchid Sarcochilus falcatus	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Orange-tip Finger- orchid Caladenia aurantiaca	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Ornate Pink-fingers Caladenia ornata	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Outcrop Guinea-flower Hibbertia hermanniifolia subsp. recondita	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Oval-leaf Grevillea Grevillea miqueliana subsp. Miquelia	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Oval-leaf Pseudanthus Pseudanthus ovalifolius	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations.
Ovens Everlasting Ozothamnus stirlingii	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Pale Flax-lily Dianella longifolia var. longifolia s.l.	Population	Bendigo FMA	Minimise disturbance from road management activities.
Pink Mountain-correa Correa lawrenceana var. cordifolia	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Pinkwood Beyeria lanceolata	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is

Species Name	Value	Applicable FMAs	Management Actions
			adequately protected during timber harvesting operations.
Pinnate Goodenia Goodenia macmillanii	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Plains Joyweed Alternanthera sp. 1 (Plains)	Population	Bendigo FMA	Minimise disturbance from road management activities.
Pomonal Leek-orchid Prasophyllum subbisectum	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Poverty Wattle Acacia dawsonii	Population	Gippsland FMAs North East FMAs	
Prawn Greenhood Pterostylis pedoglossa	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Prickly Tree-fern Cyathea leichhardtiana	Population	Gippsland FMAs East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Privet Pomaderris Pomaderris ligustrina subsp. ligustrina	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Prostrate Cone-bush Isopogon prostratus	Population	Gippsland FMAs	Apply a protection area over each population based on field inspection.
Purple Blown-grass Lachnagrostis punicea subsp. punicea	Population	Bendigo FMA	Minimise disturbance from road management activities.
	Population	Midlands FMA Gippsland FMAs	Apply a protection area over each population based on field inspection.
-			

Species Name	Value	Applicable FMAs	Management Actions
			Note: Within the Midlands FMA , the Secretary intends to review this strategy when 10 sites and 1000 individuals are protected.
Purple Diuris Diuris punctata var. punctata	Population	North East FMAs East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Purple Eyebright Euphrasia collina subsp. muelleri	Population	Gippsland FMAs North East FMAs Portland- Horsham FMA	Apply a protection area of 200 m radius over each population.
	Population	Otways FMA	Manage occurrences in consultation with the Department unless already protected.
Ridged Water-milfoil Myriophyllum porcatum	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Rigid Spider-orchid Caladenia tensa	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
River Leafless Bossiaea Bossiaea riparia	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
River Swamp Wallaby- grass Amphibromus fluitans	Population	Mid-Murray FMA	Apply a management area of 100 m radius over populations that are not already protected. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Rock daisy Brachyscome petrophila	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Rock Grevillea Grevillea willisii	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Rock Orchid Thelychiton speciosus	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Rock Tussock-grass Poa petrophila	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
	Population	Gippsland FMAs North East FMAs	Apply a protection area over each population.
Rough Eyebright Euphrasia scabra	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Rough-grain Love- grass Eragrostis trachycarpa	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Royal Grevillea Grevillea victoriae s.s.	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Rusty Velvet-bush Lasiopetalum ferrugineum var. ferrugineum	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Sand Rush Juncus psammophilus	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
	Population	Bendigo FMA	Minimise disturbance from road management activities.
Sandfly Zieria Zieria smithii subsp. smithii	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Sandpaper Fig Ficus coronata	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Scaly Greenhood Pterostylis hamata	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Scented Bush-pea Pultenaea graveolens	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.
Scented Daisy-bush Olearia tenuifolia	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Selma Saddle Grevillea Grevillea miqueliana subsp. cincta	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Shingle Fireweed Senecio diaschides	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Shining Anchor Plant Discaria nitida	Population	Gippsland FMAs	Apply a protection area over each population.
Shiny Leionema Leionema lamprophyllum	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Short Spider-orchid Caladenia brachyscapa	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Short Water-starwort Callitriche brachycarpa	Population	Otways FMA	Manage occurrences in consultation with the Department unless already protected.
Short-awned Wheat- grass Anthosachne kingiana subsp. multiflora	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
·	Population	Mid-Murray FMA	Avoid disturbance to populations.
Short-bristle Wallaby- grass Rytidosperma setaceum var. brevisetum	Population	Bendigo FMA	Minimise disturbance from road management activities.
Showy Willow-herb Epilobium pallidiflorum	Occurrences	Otways FMA	Manage occurrences in consultation with the Department unless already protected.
Sickle-leaf Rush Juncus falcatus subsp. falcatus	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Sikh's Whiskers Pterostylis boormanii	Population	Bendigo FMA	Apply a management area over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Silky Glycine Glycine canescens	Population	Bendigo FMA	Minimise disturbance from road management activities.
Silver Stringybark Eucalyptus alligatrix subsp. alligatrix	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Silver Tea-tree Leptospermum multicaule	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is

Species Name	Value	Applicable FMAs	Management Actions
			adequately protected during timber harvesting operations.
Skeleton Vine Clematis leptophylla	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Slender Bitter-cress Cardamine tenuifolia	Population	North East FMAs	Protect populations from disturbance where possible.
Slender Club-sedge Isolepis congrua	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Slender Darling-pea Swainsona murrayana	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Slender Gingidia Gingidia harveyana	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Slender Leek-orchid Prasophyllum parviflorum	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Slender Love-grass Eragrostis exigua	Population	North East FMAs	Apply a protection area of 200 m radius over each population.
Slender Mud-grass Pseudoraphis paradoxa	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Slender Parrot-pea Almaleea capitata	Population	Gippsland FMAs North East FMAs East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Slender Pomaderris Pomaderris phylicifolia	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Slender Ruddyhood Pterostylis aciculiformis	Population	North East FMAs	Protect populations from disturbance where possible.
Slender Saw-sedge Gahnia microstachya	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Slender Sword-sedge Lepidosperma gunnii	Population	North East FMAs	Protect populations from disturbance where possible.
	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Slender Tick-trefoil Desmodium varians	Population	North East FMAs	Protect populations from disturbance where possible.
-	Population	Bendigo FMA	Minimise disturbance from road management activities.
-	Population	Mid-Murray FMA	Avoid disturbance to populations.
Slender Tree-fern Cyathea cunninghamii	Population	Gippsland FMAs East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
_	Population	Otways FMA	Manage occurrences in consultation with the Department unless already protected.
Slender Violet-bush Hybanthus monopetalus	Population	Gippsland FMAs North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Small Chocolate-lily Arthropodium sp. 3 (aff. strictum)	Population	North East FMAs	Protect populations from disturbance where possible.
Small Milkwort Comesperma polygaloides	Site	Bendigo FMA	Prevent use of machinery or vehicles on sites when soils are wet or such use will damage the underground parts of the plants.
Small Scurf-pea Cullen parvum	Population	Mid-Murray FMA	Apply a management area of 100 m radius over populations that are not already protected. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Small-leaf Bush-pea Pultenaea foliolosa	Population	Gippsland FMAs North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Smooth Wallaby-grass Rytidosperma laeve	Population	Mid-Murray FMA	Avoid disturbance to populations.
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Snow Aciphyll Aciphylla glacialis	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Snow Beard-heath Acrothamnus montanus	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Snow Coprosma Coprosma nivalis	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Snow Fescue Hookerochloa eriopoda	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.

Species Name	Value	Applicable FMAs	Management Actions
Snow Pennywort Diplaspis nivis	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Snow Wallaby-grass Rytidosperma nivicola	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Snowdrop Wood-sorrel Oxalis magellanica	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Snowy River Wattle Acacia boormanii	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Soft Ledge Grass Poa hothamensis var. Parviflora	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Southern Pipewort Eriocaulon australasicum	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Spicy Everlasting Ozothamnus argophyllus	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Spinning Gum Eucalyptus perriniana	Population	Gippsland FMAs North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Spiral Sun-orchid Thelymitra matthewsii	Population	Gippsland FMAs Portland- Horsham FMA East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Sprawling Cassia Senna aciphylla	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Spreading Clubmoss Lycopodium scariosum	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Spreading Eutaxia Eutaxia microphylla var. diffusa	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.
Spreading Knawel Scleranthus fasciculatus	Population	Gippsland FMAs North East FMAs	
Squat Picris Picris squarrosa	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.
Stalked Adder's-tongue Ophioglossum reticulatum	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Stalked Brooklime Gratiola pedunculata	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Stalked Guinea-flower Hibbertia pedunculata s.l.	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Star Cucumber Sicyos australis	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Sticky Wattle Acacia howittii	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Stiff Groundsel Senecio behrianus	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Stony Bush-pea Pultenaea lapidosa	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Strawberry Buttercup Ranunculus collinus	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Striped Pink-fingers Caladenia carnea var. subulata	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Striped Pomaderris Pomaderris pilifera subsp. pilifera	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Strzelecki Gum Eucalyptus strzeleckii	Population	Gippsland FMAs	Apply a protection area over each population.
Sturdy Leek-orchid Prasophyllum aff. validum B	Population	Bendigo FMA	Apply a management area over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .

Species Name	Value	Applicable FMAs	Management Actions	
Subalpine Baeckea Baeckea latifolia	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.	
Summer Fringe-sedge Fimbristylis aestivalis	Population	Mid-Murray FMA	Avoid disturbance to populations.	
Summer Leek-orchid Prasophyllum uvidulum	Population	North East FMAs	Apply a protection area of 200 m radius over each population.	
Swamp Bush-pea Pultenaea weindorferi	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.	
Swamp Everlasting Xerochrysum palustre	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.	
Swamp Fern Thelypteris confluens	Population North E		As Apply a protection area of 200 m radius ove each population.	
Swamp Fireweed Senecio psilocarpus	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.	
Swamp Greenhood Pterostylis tenuissima	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.	
Tailed Eyebright Euphrasia caudataPopulationGippsland FMAs North East FMAs		Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.		
Tall Acrotriche Acrotriche leucocarpa	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.	
Tall Astelia Astelia australiana	Tall Astelia colony	Central Highlands FMAs	Apply a 100 m buffer around each Tall Astelia colony. Discovery of a previously unknown colony during harvesting operations will not necessarily cause harvesting to cease, but every reasonable effort will be made to protect it. Where Tall Astelia occurs within Cool Temperate Rainforest or Riparian Thicket EVCs, apply a 100 m buffer around the EVC. Buffer widths may be smaller in cases where existing roads or ridgelines occur within the 100 m. Upstream of Tall Astelia colonies, apply a minimum vegetated buffer of 40 m between the valley floor, or 'wetted zone' and adjacent	

Species Name	Value	Applicable FMAs	Management Actions	
			logging coupes. Where there is no discernible valley floor extend the buffer zone 40 m from the creek itself. Apply a 40 m buffer around isolated plants growing on road batters. No roads or snig tracks may be constructed through Tall Astelia colonies. Avoid constructing roads or snig tracks through Tall Astelia buffers. Any roads or snig tracks that do cross Tall Astelia buffers must be located as far upstream from the Tall Astelia colony as possible, and sedimentation of the Tall Astelia colony must be prevented. Where Tall Astelia occurs on seepage zones immediately below a coupe, snig tracks must be designed to minimise alteration to the sites normal drainage patterns.	
	Tall Astelia colony	Otways FMA	Apply a 100 m buffer around each Tall Astelia colony. Discovery of a previously unknown colony during harvesting operations will not necessarily cause harvesting to cease, but every reasonable effort will be made to protect it.	
Tall Vanilla-lily Arthropodium sp. 1 (robust glaucous)	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.	
Tangled Pseudanthus Pseudanthus orbicularis	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.	
Tasmanian Bladderwort Utricularia monanthos	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.	
Tasmanian Wax-flower Philotheca virgata	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.	
Tawny Spider-orchid Caladenia fulva	Population	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.	
Thick Bent-grass Deyeuxia crassiuscula	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.	

Species Name	Value	Applicable FMAs	Management Actions
Thick Eyebright Euprasia crassiuscula	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Thick-lip Spider-orchid Caladenia tessellata	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Tick Indigo Indigofera adesmiifolia	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Tight Bedstraw Galium curvihirtum	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Tingaringy Gum Eucalyptus glaucescens	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Tiny Arrowgrass Triglochin minutissima	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Tiny Bent Agrostis australiensis	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Tiny Daisy Brachyscome ptychocarpa	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions
Tiny Midge-orchid Corunastylis nuda	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Tiny Spyridium Spyridium cinereum	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Toothed leionema Leionema bilobum subsp. serrulatum	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Trailing Hop-bush Dodonaea procumbens	Population	Portland- Horsham FMA Bendigo FMA	Apply a management area over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations .
Tree Geebung Persoonia arborea	Individual tree	Central Highlands FMAs	Protect individual trees with a DBHOB of at least 10 cm from disturbance where reasonably practicable.
Tufted Club-sedge Isolepis wakefieldiana	Occurrence	Otways FMA	Manage occurrences in consultation with the Department unless already protected.
Tullach Ard Grevillea Grevillea polychroma	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Turnip Copperburr Sclerolaena napiformis	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Tussock Sedge Carex iynx	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Tussock Sedge Carex iynx	Population	North East FMAs	Protect populations from disturbance where possible.

Species Name	Value	Applicable FMAs	Management Actions
Tussock Woodrush Luzula alpestris	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Two-colour Panic Panicum simile	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Umbrella Grass Digitaria divaricatissima var. divaricatissima	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Upright Panic Entolasia stricta	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Upright Pomaderris Pomaderris virgata	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Veined Beard-heath Leucopogon neurophyllus	Population	Midlands FMA	Apply a protection area over each population.
Velvet Apple-berry Billardiera scandens s.s.	Population	Gippsland FMAs North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Velvety Geebung Persoonia subvelutinana	Population	Gippsland FMAs North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Victorian Flat-pea Platylobium alternifolium	Population	Midlands FMA	Avoid disturbance to populations within the management area, SMZ and GMZ.

Species Name	Value	Applicable FMAs	Management Actions
Wallaby-bush Beyeria lasiocarpa	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Warby Range Swamp- gum Eucalyptus cadens	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the population is adequately protected during timber harvesting operations.
Wedge Diuris Diuris dendrobioides	Population	North East FMAs	Apply a protection area of 200 m radius over each population.
Wedge Guinea-flower Hibbertia diffusa	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Wedge Oschatzia Oschatzia cuneifolia	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Western Water- starwort Callitriche cyclocarpa	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Wetland Blown-grass Lachnagrostis palustris	Population	North East FMAs	Protect populations from disturbance where possible.
	Population	North East FMAs	Protect populations from disturbance where possible.
White Billy-buttons Craspedia alba	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Whiteroot Lobelia purpurascens	Occurrence	Otways FMA	Manage occurrences in consultation with the Department unless already protected.

Species Name	Value	Applicable FMAs	Management Actions
Wild Sorghum Sorghum leiocladum	Population	Gippsland FMAs East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Wimmera Spider- orchid Caladenia lowanensis	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Winged Everlasting Ammobium alatum	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Winged Peppercress Lepidium monoplocoides	Population	Mid-Murray FMA	Apply a management area of 100 m radius over populations that are not already protected. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Winged Water-starwort Callitriche umbonata	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Wiry Wallaby-grass Plinthanthesis paradoxa	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Wombargo Wattle Acacia tabula	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Woodland Leek-orchid Prasophyllum aff. validum B	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.

Species Name	Value	Applicable FMAs	Management Actions
	Population	Mid-Murray FMA	Avoid disturbance to populations.
Woolly Buttons Leiocarpa panaetioides	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Woolly-bear Wattle Acacia lucasii	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Woolly-head Pomaderris Pomaderris eriocephala	Population	Gippsland FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Woolly Wattle Acacia lanigera var. lanigera	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.
Wrinkled Buttons Leiocarpa gatesii	Occurrence	Otways FMA	Manage occurrences in consultation with the Department unless already protected.
Wrinkled Cassinia Cassinia rugata	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.
Wrinkle-nut Lignum <i>Muehlenbeckia rhyticarya</i>	Population	Gippsland FMAs	Apply a protection area of 200 m radius over each population.
Yarra Gum Eucalyptus yarraensis	Individual tree	Midlands FMA	Retain all individual Yarra Gum trees.
Yellow Elderberry Sambucus australasica	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.

Species Name	Value	Applicable FMAs	Management Actions	
Yellow Flat-sedge Cyperus flavidus	Population	North East FMAs	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.	
	Population	North East FMAs	Apply a protection area over each population.	
Yellow Hyacinth-orchid — Dipodium hamiltonianum	Population	East Gippsland FMA	Apply a management area of 200 m radius over populations. Conduct a site inspection and detailed planning in consultation with the Department to ensure the species is adequately protected during timber harvesting operations.	
Yellow-lip Spider- orchid Caladenia xanthochila	Occurrence	Portland- Horsham FMA	Manage occurrences in consultation with the Department unless already protected.	

Table 15 Detection based rules for Heathland, Snow Gum, Swamp Gum and Heath vegetation communities

FMA	Common name	Scientific name	Management actions
East Gippsland	Heathland (all heathland EVCs and other treeless vegetation)		Apply a protection area over full extent of heathland and other treeless vegetation communities.
Midlands	Snow Gum	Eucalyptus pauciflora	Apply a protection area over full extent of vegetation communities dominated by Snow Gum on Mt Cole.
Midlands	Swamp Gum	Eucalyptus ovata	Apply a protection area over full extent of vegetation communities dominated by Swamp Gum near Enfield.
Otway	Heathy Woodland, Wet Heath and Coastal Complex		Apply a protection area over all examples of these vegetation communities.

Table 16 Buffer widths for Rainforest Sites of Significance by category and priority

Site of Significance category	Priority			
Site of Significance category	1	2	3	4
National	Extend to nearest watershed boundary	100 m	60 m	60 m
State	60 m	60 m	40 m	40 m
Regional	40 m	40 m	40 m	40 m

Table 17 Landscape management prescriptions

Applicable FMAs	Туре	Place
Central Highlands FMAs	Designated Lookout	Environs of the Mount Toorongo lookout: Prominent views of Mt. Toorongo.
Central Highlands FMAs	Designated Lookout	Morris Lookout: Lookout
Central Highlands FMAs	Designated Lookout	Natural environs of the Mounts Despair and Disappointment lookouts: Mt Disappointment summit.
Central Highlands FMAs	Designated Lookout	Prominent views from Neerim: Tarago River SMZ and North Hells Gate lookout.
Central Highlands FMAs	Designated Lookout	Seven Acre Rock: Prominent views of Blue Range.
Central Highlands FMAs	Designated Lookout	Spion Kopje Lookout: 50 m either side of the walking track and lookout.
Central Highlands FMAs	Landscape Feature	Alpine Walking Track: 50 m either side of Alpine Walking Track.
Central Highlands FMAs	Landscape Feature	Caringal Scout Camp:
Central Highlands FMAs	Landscape Feature	Dom Dom Saddle: SMZ representing prominent views of this place.
Central Highlands FMAs	Landscape Feature	Enoch's Point township
Central Highlands FMAs	Landscape Feature	Forested escarpments (Black Range and Blue Range) as viewed from the Maroondah and Goulburn Valley Highways: SMZ representing prominent views of Black Range and Blue Range.
Central Highlands FMAs	Landscape Feature	Labertouche Caves:
Central Highlands FMAs	Landscape Feature	Landscape values of the land bordering the Corranderrk and O'Shannassy aqueducts: Prominent views of Mt. Toole-be-wong .
Central Highlands FMAs	Landscape Feature	Little Yarra Valley: Prominent views of Britannia Range.
Central Highlands FMAs	Landscape Feature	Matlock Mining Area township
Central Highlands FMAs	Landscape Feature	Mt St Leonard: SMZ representing prominent views of this place.
Central Highlands FMAs	Landscape Feature	Murchison Fall and Strath Falls and their environs:
Central Highlands FMAs	Landscape Feature	Natural environs of Mount Beenak: Prominent views from Little Yarra Road.
Central Highlands FMAs	Landscape Feature	Opposite Rawson Caravan park: 50 m wide area opposite Rawson Caravan park.
Central Highlands FMAs	Landscape Feature	South Hells Gate: Prominent views from Neerim.
Central Highlands FMAs	Landscape Feature	Sylvia Creek falls and its environs

Applicable FMAs	Туре	Place
Central Highlands FMAs	Landscape Feature	Views from Mt Donna Buang and from the Warburton Highway: 100 m along the Warburton Highway. Including prominent views from Mt Donna Buang and in particular of Mt Little Joe and Mt Tugwell.
Central Highlands FMAs	Landscape Feature	Warburton-Woods Point Road / Reefton Spur Road and Upper Yarra Reservoir Rd: 50 m either side of Warburton-Woods Point Road / Reefton Spur Road and Upper Yarra Reservoir Rd. Include prominent views of Cement Creek, Armstrong Creek.
Central Highlands FMAs	Scenic Drive	Acheron Way: 50 m either side of Acheron Way.
Central Highlands FMAs	Scenic Drive	Baw Baw Road: 50 m either side of Baw Baw Road.
Central Highlands FMAs	Scenic Drive	Eildon Road: 50 m either side of Eildon Road.
Central Highlands FMAs	Scenic Drive	Forested escarpments of the Mounts Tallarook and Disappointment forest as viewed from the Hume Highway: Prominent views of Mt Disappointment.
Central Highlands FMAs	Scenic Drive	Landscape of Pauls Range: 50 m either side of Chum Creek Road.
Central Highlands FMAs	Scenic Drive	Maroondah Highway: 100 m either side of Maroondah highway, prominent views.
Central Highlands FMAs	Scenic Drive	Marysville - Buxton Road: 50 m either side of Marysville - Buxton Road, and prominent views of Mt. Strickland, Triangle Block and Mt. Margaret.
Central Highlands FMAs	Scenic Drive	Melba Highway: 100 m either side of Melba Highway.
Central Highlands FMAs	Scenic Drive	Moe - Walhalla Road: 50 m either side of Moe - Walhalla Road.
Central Highlands FMAs	Scenic Drive	Murrindindi Road from Murrindindi to Murrindindi Cascades: 50 m either side of Murrindindi Road from Murrindindi to Murrindindi Cascades.
Central Highlands FMAs	Scenic Drive	Myers Creek Road: 50 m either side of Myers Creek Road.
Central Highlands FMAs	Scenic Drive	Myers Creek Scenic Reserve: 50 m either side of Myers Creek Road.
Central Highlands FMAs	Scenic Drive	Natural environs of Lady Talbot Drive, Marysville: 50 m either side of Lady Talbot Drive.
Central Highlands FMAs	Scenic Drive	Powelltown - Noojee Road: 50 m either side of Powelltown - Noojee Road.
Central Highlands FMAs	Scenic Drive	Powelltown - Noojee Road: 50 m either side of Powlltown - Noojee Road.
Central Highlands FMAs	Scenic Drive	Steavenson Falls Road: 50 m either side of Steavenson Falls Road.
Central Highlands FMAs	Scenic Drive	Thomson Valley - Mt St Gwinear Rd: 50 m either side of Thomson Valley - Mt St Gwinear Rd.
Central Highlands FMAs	Scenic Drive	Thomson Valley Road from Erica to the Thomson Reservoir : 50 m either side of Thomson Valley Road from Erica to the Thomson Reservoir. Include prominent views of Mt. Erica and Erica surrounds.
Central Highlands FMAs	Scenic Drive	Whittlesea - Yea Road: 50 m either side of Whittlesea - Yea Road.

Applicable FMAs	Туре	Place
East Gippsland	Designated Lookout	Delegate Hill
East Gippsland	Designated Lookout	Genoa Peak
East Gippsland	Designated Lookout	Gunmark lookout
East Gippsland	Designated Lookout	Maramingo Hill
East Gippsland	Designated Lookout	Mount Bemm
East	Designated	Mount Bowen
Gippsland East	Lookout Designated	Mount Buck
Gippsland East	Lookout Designated	Mount Denmarsh
Gippsland East	Lookout Designated	Mount Ellery
Gippsland East	Lookout Designated	<u> </u>
Gippsland East	Lookout Designated	Mount Kaye
Gippsland East	Lookout Designated	Mount McLeod
Gippsland	Lookout	Mount Nowa Nowa
East Gippsland	Designated Lookout	Mount Raymond
East Gippsland	Designated Lookout	Mount Seldom Seen
East Gippsland	Designated Lookout	Mount Stewart
East Gippsland	Designated Lookout	Mount Tingaringy
East Gippsland	Designated Lookout	Mount Tower
East Gippsland	Landscape Feature	Buchan: Black Satin Creek, Site # 810/07
East Gippsland	Landscape Feature	Buchan: Wattle Gully, Site # 812/09
East	Landscape	Buchan: W-Tree Creek headwaters, Site # 813/06
Gippsland East	Feature Scenic	Betka Forest Drive: Mallacoota and Croajingalong access; native forest
Gippsland East	Drive Scenic	management. Big River Road (part): Mount Ellery access.
Gippsland East	Drive Scenic	Bonang - Bendoc Road and Playgrounds Track: Day trips around Bendoc,
Gippsland East Gippsland	Scenic Drive	Gold Mining, park access. Bonang Road (Orbost - NSW border): Mt Buck/Cooney Ridge; Mt Watt; Mt Rich; views from Goongerah to Postmans Spur and BA ridge (Mt Ellery in background); Mt Little Bill; Brown Mountain; views from Bonang area to Mt Koolabra and Cottonwood Range; enclosed forest canopy and high quality roadside scenery along Pinch Swamp Ck.
East Gippsland	Scenic Drive	Buchan Road (Bruthen - Buchan): Views west from Red Knob area; Pretty Sally; vegetation between roadside and SEC easement provides variety and reduces impact of easement; Fluke Knob; Spencer Knob; Lookout Rocks; Tara Range.
East Gippsland	Scenic Drive	Buchan—Jindabyne Road: National Estate (Road built in Great Depression). Mt Dawson; views from W-Tree to Dawson range (including Half Moon Gully, Mt Elephant, Lookout Top, Gum Top and Wagtail Ridge) and Mt Murrindal; Mt Statham; Mt Seldom Seen.
East Gippsland	Scenic Drive	Cabbage Tree - Cape Conran Road, Marlo - Cabbage Tree Road and Palms Track: National Estate(Baldwin - Spencer journey); access to Cabbage Palms Flora Reserve.
East Gippsland	Scenic Drive	Cann Valley Highway: Extensive views west across flats to forested range including Mt Noorinbee, High Peak, Cann Mountain, Morgan Hill and Bennet Knob, Mt Petterson.

Applicable FMAs	Туре	Place						
East Gippsland	Scenic Drive	Clarkeville Road: Errinundra National Park access.						
East	Scenic Drive	Coast Range Road (part): Part of tall forest experience in Errinundra						
Gippsland East	Scenic	National Park; nocturnal birds and arboreal mammals. Colquhoun Forest Drive: Koalas; historic features associated with the						
Gippsland	Drive	Orbost - Bairnsdale railway. Combienbar and Errinundra Rds: National Estate (Baldwin-Spencer						
East Gippsland	Scenic Drive	journey); Errinundra National Park access; (through to Gap Rd) hills around Club Terrace; Pyramid Hill; Shining Gums and Cool Temperate Rainforest around Kanuka Creek; Tommy Roundhead Hill.						
East Gippsland	Scenic Drive	Deddick River Road (part) 36-mile Road: Snowy River National Park access.						
East Gippsland	Scenic Drive	Dellicknora / Cameron Roads: Tingaringy National Park access.						
East Gippsland	Scenic Drive	Gap Road: National Estate (Baldwin-Spencer journey); high-quality foreground scenery.						
East	Scenic	Greens Road: Access to Errinundra National Park and St Patricks Falls;						
Gippsland East	Drive Scenic	native forest management. Gunmark Road: Major access to Errinundra National Park; extensive views across Delegate River headwaters; buffers required where road forms						
Gippsland	Drive	border between the Park and State forest.						
East Gippsland	Scenic Drive	Harris Creek/ 2-mile Road: Mount Nowa Nowa access.						
East Gippsland	Scenic Drive	Hensleigh Creek Road (part): Errinundra National Park access; nocturnal birds and arboreal mammals.						
East Gippsland	Scenic Drive	Jones Creek Road: Coopracambra National Park access.						
East Gippsland	Scenic Drive	Lake Tyers House Road: National Estate (Old Coach Road); Lake Tyers State Park access.						
East Gippsland	Scenic Drive	Mallacoota and Betka Roads: Access to Mallacoota, beaches and Croajingolong National Park.						
East Gippsland	Scenic Drive	Murrungowar Forest Drive (Princes Hwy - Murrungowar Rd - Glen Arte Rd - Bendoc Ridge Rd - Princes Hwy): Jungle Hill; Glen Arte Flora Reserve; regrowth thinning; side trips to Cabbage Tree Falls and St Patricks Falls; historic features around Murrungowar and Glen Arte.						
East Gippsland	Scenic Drive	Nowa Nowa—Buchan Road: "The Gorge"; Mount Nowa Nowa; scenic alternative route to Buchan.						
East Gippsland	Scenic Drive	Nunnett and Timbarra Roads: Alpine National Park access.						
East Gippsland	Scenic Drive	Orbost-Buchan Road: National Estate (aesthetic quality).						
East Gippsland	Scenic Drive	Princes Highway (Lakes Entrance—Orbost): Enclosed forest canopy between Nowa Nowa and Lakes Entrance; stream crossings at Hospital Ck, Hartland River and Simpson's Ck.						
East Gippsland	Scenic Drive	Princes Highway (Orbost-NSW border): Stream crossings at Brodribb River (rainforest) and Jungle Ck; forested hills seen from Cabbage Tree Ck; thinning operations between Cabbage Tree and Bemm River; enclosed forest canopy between Bemm River and Lind National Park; Mt Bemm; Donalds Knob; hills to south of Tonghi Creek; views north up the Cann Valley; views north-west from Mt Drummer.						
East Gippsland	Scenic Drive	Seldom Seen Tower Road : Access to Mount Seldom Seen.						
East Gippsland	Scenic Drive	Sydenham Inlet Road: Wide clearing maintained as strategic fire-break; view from township and inlet north to hills above township.						
East Gippsland	Scenic Drive	Tamboon and Point Hicks Roads: Croajingolong National Park access; middle ground views near Furnell Landing.						
East Gippsland	Scenic Drive	Tulloch Ard Road: Snowy River National Park access.						
East	Scenic	Wallagaraugh Road: Croajingolong National Park access.						
Gippsland East	Drive Scenic	Wallagaraugh Road: Croajingolong National Park access.						
Gippsland East	Drive Scenic	Wangarabell Road: Access to Maramingo Hill; Genoa river access.						
Gippsland	Drive	vvangarabeli Koau. Access to iviaramingo filli, Genoa river access.						

Applicable FMAs	Туре	Place						
East Gippsland	Scenic Drive	West Wingan Road: Croajingolong National Park access.						
East Gippsland	Scenic Drive	Yalmy/Rising Sun/Pinnak Roads: Major access road to Snowy River National Park; forest vistas into catchments of Cavender Ck, Serpentine Ck, Yalmy River and Stony Ck.						
Gippsland FMAs	Designated Lookout	Bindi Lookout						
Gippsland FMAs	Designated Lookout	Elphick Lookout: Lookout						
Gippsland FMAs	Designated Lookout	Pendergast Lookout: Lookout						
Gippsland FMAs	Designated Lookout	Stutterin' Fred's Lookout: Lookout						
Gippsland FMAs	Landscape Feature	Blue Pool: Either side of the Freestone Creek.						
Gippsland FMAs	Landscape Feature	Buenba Flat						
Gippsland FMAs	Landscape Feature	Great Alpine Road: Prominent views.						
Gippsland FMAs	Landscape Feature	Haunted Stream Valley/Stirling Ghost Town: 300 m either side of the river						
Gippsland FMAs	Landscape Feature	Haunted Stream Valley: 300 m either side of the river.						
Gippsland FMAs	Landscape Feature	Nicholson River: 100 m downstream of Marthavale, including up to track on west side of river and 50 m west side of Nicholson track.						
Gippsland FMAs	Landscape Feature	The Brothers						
Gippsland FMAs	Scenic Drive	Boola Road: 20 m either side of the road.						
Gippsland FMAs	Scenic Drive	Bruthen – Buchan Road: 50 m either side of the road.						
Gippsland FMAs	Scenic Drive	Dargo High Plains Road: 100 m either side of the road.						
Gippsland FMAs	Scenic Drive	Deptford Road: 50 m either side of the road.						
Gippsland FMAs	Scenic Drive	Fairy Dell: 50 m either side of the associated forest drive.						
Gippsland FMAs	Scenic Drive	Great Alpine Road (part): Prominent views.						
Gippsland FMAs	Scenic Drive	Heyfield – Licola Road						
Gippsland FMAs	Scenic Drive	Hyland Highway: 50 m either side of the road.						
Gippsland FMAs	Scenic Drive	South Gippsland Highway: Prominent views .						
Gippsland FMAs	Scenic Drive	Strzelecki Highway: 50 m either side of the road.						
Mid Murray	Landscape Feature	River Track: 20 m adjacent to the whole length of the road.						
Mid Murray	Scenic Drive	Iron Punt Track: 20 m adjacent to the whole length of the road.						
Mid Murray	Scenic Drive	Lock Road: 20 m adjacent to the whole length of the road.						
Mid Murray	Scenic Drive	Nursery Track: 20 m adjacent to the whole length of the road.						
Mid Murray	Scenic Drive	Rifle Butts Track to Five Sleepers Track to Thompson Track: 20 m adjacent to the whole length of the road.						
Mid Murray	Scenic Drive	Stanton Break: 20 m adjacent to the whole length of the road.						
Midlands	Designated Lookout	Ben Nevis Picnic Area						
Midlands	Designated Lookout	Black Hill Lookout						
Midlands	Designated Lookout	Daylesford Botanical Gardens Lookout						

Applicable FMAs	Туре	Place						
Midlands	Designated Lookout	Jackson Lookout (Hepburn)						
Midlands	Designated Lookout	Lookout Hill Picnic Area						
Midlands	Designated Lookout	Mount Buangor Picnic Area						
Midlands	Designated Lookout	Sovereign Hill Lookout						
Midlands	Scenic Drive	Ararat - St Arnaud Road (Crowlands - Landsborough)						
Midlands	Scenic Drive	Bacchus Marsh - Gisborne Road						
Midlands	Scenic Drive	Ballan - Daylesford Road						
Midlands	Scenic Drive	Ballan Daylesford Road						
Midlands	Scenic Drive	Ballarat - Buninyong Road (Sovereign Hill - Buninyong)						
Midlands	Scenic Drive	Beaufort - Lexton Road						
Midlands	Scenic Drive	Bungaree Creswick Road						
Midlands	Scenic Drive	Clunes Creswick Road						
Midlands	Scenic Drive	Creswick Dean Road						
Midlands	Scenic Drive	Daylesford - Ballarat Road						
Midlands	Scenic Drive	Daylesford - Malmsbury Road						
Midlands	Scenic Drive	Daylesford - Trentham Road						
Midlands	Scenic Drive	Gisborne - Melton Road (Pyrites forest)						
Midlands	Scenic Drive	Glenelg Highway (Linton - Smythesdale)						
Midlands	Scenic Drive	Greendale - Myrniong Road						
Midlands	Scenic Drive	Greendale - Trentham Road						
Midlands	Scenic Drive	Kyneton - Trentham Road						
Midlands	Scenic Drive	Lal Lal Falls Road						
Midlands	Scenic Drive	Lerderderg River (Heritage River Section)						
Midlands	Scenic Drive	Midland Highway (Ballarat - Springmount)						
Midlands	Scenic Drive	Midland Highway (Kennedys Gully - Spingmount)						
Midlands	Scenic Drive	Midlands Highway (Clarendon - Meredith)						
Midlands	Scenic Drive	Mount Cole Main Road						
Midlands	Scenic Drive	Myrniong Trentham Road						
Midlands	Scenic Drive	Navigators Road						
Midlands	Scenic Drive	Pyrenees Highway (Avoca - St Arnaud - Ararat Road)						
Midlands	Scenic Drive	Sebastopol - Smythesdale Road (Smythesdale - Ross Creek)						
Midlands	Scenic Drive	Springs Road						

Applicable FMAs	Туре	Place					
Midlands	Scenic Drive	Sunraysia Highway (Avoca - Tanwood)					
Midlands	Scenic Drive	Western Freeway (Bacchus Marsh - Bungaree)					
Midlands	Scenic Drive	Western Highway (Mt. Langi Ghiran - Beaufort)					
Midlands	Scenic Drive	White Swan Road					
North East FMAs	Designated Lookout	Mt Murramurrangbong and Mt Big Ben					
North East FMAs	Landscape Feature	15 Mile Creek: Visual corridor and natural water feature. 50 m either side of creek.					
North East FMAs	Landscape Feature	Beechworth Forest Drive: Scenic corridor for tourist roads. 50 m each side of road.					
North East FMAs	Landscape Feature	Big Hill/The Springs Saddle/Bald Hill/Bogong Jack Saddle: Mountain views from Mt Beauty township and Mt Fainter North. Scenic corridor for tourist roads.					
North East FMAs	Landscape Feature	Bindaree Falls: 100 m radius over the falls.					
North East FMAs	Landscape Feature	Buckland Valley: Dingo Ck to Buckland River East Branch, 50 m each side of stream.					
North East FMAs	Landscape Feature	Carters Road Picnic Area: High scenic values in recreation area. 50 m radius over the picnic area.					
North East FMAs	Landscape Feature	Clear Creek (Tributary of Buckland River): 50 m each side of stream.					
North East FMAs	Landscape Feature	Conic Range (partly State forest)					
North East FMAs	Landscape Feature	Craig Hut: 200 m radius over the hut.					
North East FMAs	Landscape Feature	Dandongadale River: Visual corridor and natural water feature. 50 m each side of river.					
North East FMAs	Landscape Feature	Granite Peak: RNE Natural site. Mountain Views from Omeo Hwy. State forest above 1100 m elevation.					
North East FMAs	Landscape Feature	Harkers Creek (Mitta Mitta): 50 m each side of stream					
North East FMAs	Landscape Feature	Harrietville Valley (vicinity of Harrietville) and Ovens River, West Branch: 20 m each side of the Great Alpine Road.					
North East FMAs	Landscape Feature	Holland Creek (part): Visual corridor and natural water feature. 50 m either side of creek.					
North East	Landscape	Lightning Ridge (Strathbogie Ranges): Mountain views from Lake					
FMAs North East	Feature Landscape	Nillahcootie and Midland Highway. Mitta Mitta Valley (downstream of Mitta Mitta): NTA(V) classified, Visual					
FMAs North East	Feature Landscape	corridor and natural water feature. 50 m each side of road. Moonee Range (Strathbogie Ranges): Mountain view from Midland					
FMAs North East	Feature Landscape	Highway Mt Pinnibar					
FMAs North East	Feature Landscape	Myrtle (Barwidgee) Creek Valley (vicinity of Myrtleford): 50 m on State forest					
FMAs North East	Feature Landscape	side of Stanley to Barwidgee Road. Pheasant Creek (Tributary of Buckland River): Visual corridor and natural					
FMAs	Feature	water feature. 50 m each side of road.					
North East FMAs	Landscape Feature	Plain Creek Tramway (Sawmill Settlement): Historic site of local significance. 20 m buffer around the historic trestle bridge.					
North East FMAs	Landscape Feature	Sandy Creek Waterfall (Strathbogie Ranges): 100 m radius over the falls.					
North East FMAs	Landscape Feature	Snowy Creek Waterfall: 100 m radius over the falls.					
North East FMAs	Landscape Feature	The Pinnacles & Mt Jack including Barwidgee Creek: Headwaters of Barwidgee Creek.					
North East FMAs	Landscape Feature	Trappers Creek (near Mountain Creek): Visual corridor and natural water feature. 50 m each side of road.					
North East FMAs	Landscape Feature	Upper Delatite valley (vicinity of Mirimbah): 200 m each side of river.					
North East FMAs	Landscape Feature	Upper Nariel valley above Crawford Crossing: Visual corridor and natural water feature. 50 m each side of stream (where in State forest).					

Applicable FMAs	Туре	Place						
North East	Landscape	West Kiewa valley: Viewed from Mt Feathertop, indicative aesthetic						
FMAs	Feature	National Estate.						
North East FMAs	Landscape Feature	Wongungarra & Humffray Rivers & Teatree Range except ridge: Indicative Natural Landscape (National Estate). Located in National Park, and Mt Sarah Scenic Reserve.						
North East	Landscape	Yackandandah Creek (from Nine Mile Creek to State forest boundary):						
FMAs	Feature	Visual corridor and natural water feature. 50 m either side of creek.						
North East	Landscape	Yarrabulla Creek: Visual corridor and natural water feature. 50 m each side						
FMAs	Feature	of stream. Partly within Remote & Natural Area.						
North East	Landscape	Yarrabulla Creek: Visual corridor and natural water feature. 50 m each side						
FMAs	Feature	of stream. Partly within Remote & Natural Area.						
North East	Scenic	Nariel-Benambra Road: Scenic corridor for tourist road. 50 m each side of						
FMAs	Drive	road.						

Table 18 Recreation sites

FMA	Locality	Place	Extent of SMZ		
Gippsland FMAs	Heyfield	12 Mile Creek	50 m radius over the site.		
Gippsland FMAs	Heyfield	Gows Hotel	50 m radius over the site.		
Gippsland FMAs	Heyfield	Kinleys Yards	50 m radius over the site.		
Gippsland FMAs	Heyfield	Noon Road	50 m radius over the site.		
Gippsland FMAs	Swifts Creek	Washington Winch	200 m radius over the site.		
North East FMAs	Beechworth	Beechworth Forest Drive (Includes Clarkes Corner, Murmungee Lookout and Bates Dam en route)	Forest drive and environs.		
North East FMAs	Myrtleford	Dandongadale River	50 m radius over the site.		

Table 19 Research sites

FMA	Locality	Research project	Management Actions
Gippsland FMAs	Harraps Highway	Regrowth of Phytophthora cinnamomi dieback	Limit activities to those that are compatible with the experiment objectives.
Gippsland FMAs	Old Rosedale Road	Planting on dieback affected sites	Limit activities to those that are compatible with the experiment objectives.
Gippsland FMAs		Boola Silvertop thinning trials	Limit activities to those that are compatible with the experiment objectives.
Gippsland FMAs		Mullungdung Phytophthora cinnamomi progeny trial EUC475	Limit activities to those that are compatible with the experiment objectives.
Gippsland FMAs	Bentley's Plain Road	E. delegatensis provenance trial	Limit activities to those that are compatible with the experiment objectives.
Gippsland FMAs	Engineers Road, Barmouth Spur Track, Clarks Track, Angora Range Road	Gippsland mixed species growth study	Limit activities to those that are compatible with the experiment objectives.
Midlands	Wombat and Mt Cole SF	Armillaria research	exclude timber harvesting operations and other management activities which will affect research objectives until 2020.
Midlands	Wombat SF	Fire effects study area	exclude timber harvesting operations and other management activities until such time as the areas are no longer required.

FMA	Locality	Research project	Management Actions
Midlands	Wombat SF	Regrowth thinning and fertiliser trials	Restrict timber harvesting operations and other management operations to those that are compatible with the objectives of the research until 2020.
Midlands		Forest growth plots	Exclude timber harvesting operations and other management activities until such time as the areas are no longer required.
Midlands		Continuous Forest Inventory plots	Protect unharvested plots and a 50m buffer until 2020. Review the need for further protection at this time.
North East FMAs	Clear Hills	1957 Alpine Ash thinning trials	Limit activities to those that are compatible with the experiment objectives.
North East FMAs	Flagstaff	KTRI Paterson's Curse Crown Boring Weevil research site	Limit activities to those that are compatible with the experiment objectives.
North East FMAs	Highett Ridge Eildon Mount Elliot	KTRI St John's Wort Mite research sites	Limit activities to those that are compatible with the experiment objectives.
North East FMAs	Lords Creek Mt Wills Creek / Snowy Creek Lightning Creek Track Bullhead Callaghan Creek Long Spur	KTRI Blackberry Rust Fungus research sites	Limit activities to those that are compatible with the experiment objectives.
North East FMAs	Mt Pinnibar	1928 Alpine Ash thinning trial	Limit activities to those that are compatible with the experiment objectives.
North East FMAs	Myrtleford	Cropper Creek hydrologic research project	Limit activities to those that are compatible with the experiment objectives.
North East FMAs	Old Tolmie Road, Toombullup	CFTT Tree Breeding Research provenance trials for E. delegatensis (EUC 411 1979), E. globulus (EUC413 1981) and E. nitens (EUC 426 1990)	Limit activities to those that are compatible with the experiment objectives.
North East FMAs	Riley Creek	Long-footed Potoroo research trapping grid	Manage in accordance with Long- footed Potoroo prescriptions.

Table 20 Road classification system

Road Class ¹	Туре	Service Function Description	Road Type Description
5A	Primary Road >100 ADT ²	 Provides primarily for the main traffic movements into and through a region. This includes access to high use visitor sites and forest areas Caters generally for higher travel speed, all vehicle types including large vehicles (i.e. buses and trucks). 	 All weather road predominantly 2-lane and mainly sealed. A high quality³ of service road Design speed standard of 80 – 50 km/h according to terrain Minimum carriageway width is 7 m
5B	Secondary Road 100 – 30 ADT	 Provides access to moderate use visitor sites and forest areas Serves the purpose of collecting and distributing traffic from local areas, moderate use visitor sites and forest areas to or from primary or minor roads Caters for moderate travel speed a full range of vehicles including large vehicles 	 All weather 2 lane road formed and gravelled or single lane sealed road with gravel shoulders A good quality of service road Design speed standards of at 70 – 30 km/h according to terrain
5C	Minor Road 50 – 20 ADT	 Provides a link to low and moderate use visitor sites and forest areas, and forms a feeder link to a logging coupe access track/road or fire track Purpose is to link areas, which are traffic generators to secondary or primary roads Caters for lower travel speed and full range of vehicles 	 Generally all-weather single lane 2-way unsealed formed road usually lightly gravelled A fair quality of service road Design speed standards of 60 – 20 km/h according to terrai
5D	Access Track/Road <20 ADT	 Provides access to low use visitor sites and forest areas Can be short term, temporary or feeder roads to access individual timber harvesting coupes Provides for fire protection and management access (sometimes exclusively for management vehicles only) Caters for low travel speed and a range of vehicles in dry weather May be seasonally closed 	 Substantially a single lane 2-way generally dryweather, formed (natural materials) track/road A low quality of service track/road Design speeds of 40 - <20 km/h according to terrain May be restricted to 4 wheel-drive vehicles
5E	Rough Track <10 ADT	 Provides primarily for 4 wheel-drive vehicles Mainly used for fire protection purposes, management access and limited recreational activities Caters for very low travel speed May be seasonally closed May be restricted to management vehicles only 	 Predominantly a single lane 2-way earth tracks (unformed) at or near the natural surface level A very low quality of service track Predominantly not conforming to any geometric design standards

Notes: ¹ The prefix 5 is related to the Austroads National Functional Road Classification categories.

² Average Daily Traffic (ADT) represents the average daily traffic over the peak season.

³ Quality of service is defined in terms of the level of convenience, (geometric standards), comfort (ride qualities) and safety (travel consistency) provided

Table 21 Guidelines to the main geometric design standards – unsealed roads

Main Geometric Characteristic	5B Sec	ondary		5C Mind	or		5D Acce	ess		5E Trac	ks		Comments
	Flat	Rolling	M'tain	Flat	Rollin g	M'tain	Flat	Rollin g	M'tain	Flat	Rolling	M'tain	
Speed value km/hr	70	50	30	60	40	20	40	30	<20	N/A	N/A	N/A	Based on 85th percentile speed
Cross section elements													
Number of traffic lanes	2	2	2	1	1	1	1	1	1	1	1	1	5B – two lane unsealed
Minimum cross fall/cambe	er %												
Sealed road	3	3	3	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	
Unsealed road	5	5	5	5	5	5	5	5	5	4	4	4	Min of 4% to drain rain fall off tracks
Max superelevation % (2)													
Sealed road	6-8	6-10	12	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	Actual value based on curve radius adopted
Unsealed road	6	8	10	6	8	10	6	8	10	N/A	N/A	N/A	
Min traffic lane width m (3)	3	3	3	3	3	3	3	3	3	3	3	3	Unsealed values
Min shoulder widths m	0.5	0.5	0.5	1.5	1	0.5	1	0.5	0.5	0	0	0	Unsealed values
Min carriageway width (lanes + shoulders) m	7	7	7	6	5	4	5	4	4	3	3	3	Unsealed values
Min formation width (including verges	9	9	9	8	7	6	7	6	6	3	3	3	Unsealed values
Horizontal Geometry													
Min radius curve m (5)													
Sealed road	110	45	15	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	
Unsealed road	250	100	35	170	60	15	70	35	15	N/A	N/A	N/A	
Min stopping sight distan	ce m (6)												
Sealed road	90	50	25	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	At tight curves check for lateral obstructions to

Main Geometric Characteristic	5B Sec	ondary		5C Mind	or		5D Acce	ess		5E Trac	ks		Comments
	Flat	Rolling	M'tain	Flat	Rollin g	M'tain	Flat	Rollin g	M'tain	Flat	Rolling	M'tain	
													provide the stopping sight distance required.
Unsealed road	120	70	30	90	50	30	50	30	20	N/A	N/A	N/A	
Min meeting sight dista	nce m (7)												
Sealed road	170	100	45	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	For high volume roads may need to check for overtaking provisions
Unsealed road	230	130	60	180	100	60	100	60	40	N/A	N/A	N/A	
Vertical Geometry													
Max vertical grade % (8)	6	8	12	6	8	12	6	8	15	N/A	N/A	N/A	For tracks avoid steep grades to reduce soil erosion
Min crest vertical curve K values (9)	28	9	5	9	4	2	4	2	1	N/A	N/A	N/A	Unsealed values
Min sag vertical curve K values (10)	8	4	3	6	3	2	3	2	1	N/A	N/A	N/A	Unsealed values

- (1) A simple classification of 'flat', 'rolling' and 'mountainous' terrain descriptions has been adopted as a basis for specifying appropriate geometric standards. The definition of each can be described in general terms as follows:
 - Flat terrain Roads generally follow the ground contours. Typical gradients being up to 1:20 or 5 % (<3 degrees).
 - Rolling terrain Roads can have substantial cuts and fills. Typical gradients being up to 1:20 to 1:8 or 5 12.5 % (3-7 degrees)
 - Mountainous terrain Roads in rugged hilly or mountainous ground with substantial restrictions to both horizontal and vertical alignments. Typical gradients being > 1:8 or > 12.5 % (7 degrees).
- (2) The maximum superelevation value will need to take into account the use of the road by high loaded heavy vehicles, speed and curve radii.
- (3) In cases where there is a high percentage of heavy vehicles (>20%) minimum lane widths can be increased by 0.5 m.
- (4) Allows for 1m verge/table drain width. This must be reviewed based on actual locations where for drainage reasons greater widths may be required.
- (5) This is based on Austroads (1989)and Giummarra G. (2001). Lower values of surface coefficients on unsealed roads result in radii being greater. Values rounded up. For minimum radius curves widening on the inside of a curve may be necessary to accommodate longer vehicles.
- (6) Based on reaction time of 2 seconds and surface coefficients relating to sealed and unsealed surfaces and values round up. Values based on flat grades and allowances will need to be made for up and down grades.

- (7) This is a requirement for single lane two-way roads. Values rounded up.
- (8) In some cases higher grades of up to 20% can be allowed for short sections (about 150 m). Keep grades on unsealed roads lower due to ravelling and scouring of surface.
- (9) Calculation of these values if to be based on information contained in Austroads (1989) for sealed roads and Giummarra G. (2000) for unsealed roads. The length of the vertical curve (L) is based on the product of K multiplied by the algebraic difference in grades percentage A. (i.e. L+K×A).
- (10) Sag values are based on comfort control criteria.

Table 22 Minimum clearing widths (m) required for typical road construction

Side Slope		Road Class					
Degrees	%	5B	5C	5D	5E		
0-7.5	0-13	13	7	6	4		
7.5-15	13-27	17	11	10	7		
15-22.5	27-41	23	17	16	10		
22.5-30	41-58	30	24	23	14		

Notes:

- Refer to **Table 20 Road classification system** above for road class definitions.
- The recommended clearing width is a horizontal distance.
- Table 22 Minimum clearing widths (m) required for typical road construction above assumes that debris disposal is complete before construction commences.
- Extra clearing width may be required if debris disposal is concurrent with road construction.
- Clearing widths do not include slashed verges except within engineering or **road** safety priority areas where extra cleared widths are shown to be required.
- The recommended clearing width for **temporary road** is 6m.
- Additional widening is often required at curves.

Table 23 Maximum distance between drainage structures

0-11		ROAD GRADE								
Soil Erosion Hazard	1:50 1º 2%	1:25 2º 4%	1:15 3.5° 6%	1:12 4.5° 8%	1:10 6º 10%	1:8 7º 12%	1:7 8º 15%	1:5 11º 20%		
Low	250	170	130	115	100	90	60	30		
Moderate	200	150	120	105	90	80	50	NP ¹		
High	160	130	110	95	80	65	NP ¹	NP ¹		

Notes:

- Drainage structures include culverts, roll-overs, and inverts or cross drains.
- ¹ Road Grade Not Permitted.

Table 24 Fire severity classification

Severity class	Severity type	Description
	arevus hum	90 – 100 % of eucalypt and non-eucalypt crowns are burnt.
ı	crown burn	an intense burn with widespread crown removal.
		60 – 100 % of eucalypt and non-eucalypt crowns are scorched some crowns are burnt.
2	crown scorch	
		an intense understorey fire with complete crown scorch of most eucalypt and non-eucalypts.
		30 – 65 % of eucalypt and non-eucalypt crowns are scorched.
3	moderate crown scorch	a variable intensity of fire ranging from a warm ground burn with no crown scorch to an intense understorey fire with complete crown scorch of most eucalypt and non-eucalypts.
		1 – 35 % of eucalypt and non-eucalypt crowns are scorched.
4	light crown scorch	a light ground burn with isolated patches of intense understorey fire and some crown scorch.
5	no crown scorch	< 1 % of eucalypt and non-eucalypt crowns are scorched, understorey may be burnt or unburnt.

Figure 2 Field process for determining protection areas and management areas based on modelled or mapped values




Table 25 Salvage harvesting Barred Galaxia minimum buffer and filter strip widths

Stream Class	Slope 0-20°	Slope 21-30°
Permanent	40 m B + 10 m F	50 m B + 10 m F
Temporary	20 m B + 20 m F	30 m B+ 20 m F
Drainage Lines	10 m F	15 m F
Wetlands	40 m B	50 m B

Note: Slope is the average slope of the **coupe** and potential **buffer** area situated within 100 m of the **waterway** or to the top of the drainage basin (whichever is closest).

Table 26 Basal Area (BA) and stocking specifications for commercial ash thinning

Stand age (yrs.)	Minimum retained BA (m²/ha)	Stocking range (stems/ha)	Average square spacing (m)		
14-20	17	300-400	5.3		
21-30	20	230-320	6.0		
31-35	23	170-250	6.9		
36-40	28	120-190	8.0		
41+	32	80-140	9.5		